

STUDIA GEOBOTANICA

An international journal

Vol. 11 1991

EDITORS

G. Estabrook - Ann Arbor, Mi
L. Iljanic - Zagreb
E. Mayer - Ljubljana
L. Orlóci - London, On.
F. Pedrotti - Camerino
S. Pignatti - Roma
A. Pirola - Pavia
J. Poelt - Graz
L. Poldini - Trieste
E. Wikus Pignatti - Trieste

EDITOR IN CHIEF

D. Lausi - Trieste

SECRETARY

P.L. Nimis - Trieste



**Dipartimento di Biologia
Sezione di Geobotanica ed Ecologia vegetale**

Università di Trieste

INSTRUCTIONS TO AUTHORS

STUDIA GEOBOTANICA publishes original articles in the field of geobotany. It also publishes book reviews, notices of scientific meetings and summaries of proceedings.

MANUSCRIPTS should be submitted in duplicate to an Editor or to the Secretary of the Editorial Board. The Secretary will confirm receipt of the manuscript. The suitability of submitted manuscripts will be judged by appropriate members of the Editorial Board. For specialized articles, appropriate referees will be consulted. The Secretary will inform authors of the acceptance or rejection of manuscripts or the need for substantial revisions or reduction.

Manuscript should be typed, double-spaced, on white bond paper with a 25 mm margin on all sides.

LANGUAGE should be preferably English, Italian, French and German are also accepted.

THE TITLE should be informative and brief. The name (s) of the author (s) should be placed below the title. References to research projects, institutional publication numbers, etc. should be placed in a footnote to the author's names.

Acknowledgements and dedications should be simply phrased and given at the end of the article.

KEYWORDS, no more than 10, in English, should be placed on the title page in alphabetical order.

ABSTRACT. An English abstract must be included. A summary in either French, German, Italian or in Slavic language may also be added. It should not exceed one manuscript page.

LITERATURE. For examples see the published *Studia Geobotanica*.

CHAPTER HEADINGS should be given without designations, second or third order subheadings are permitted.

TABLES should be typed on separate pages, and not in the text itself. They should preferably be submitted in camera-ready form, i.e. typed on an IBM electric typewriter (using, e.g., an 'Artisan' typing element and carbon ribbon). Tables should be typed without corrections in such a way that they can be reduced to one full page (width 120 mm). Large tables should be prepared in the same way. If they cannot be reduced to page-size, they will be inserted as fold-out tables: if possible, this should be avoided. Tables must be numbered and given descriptive titles. If the tables are not prepared according to these instructions, the publishers reserve the right to ask for a contribution to the actual cost.

ILLUSTRATIONS. Figures such as drawings and graphs should be done in India ink on smooth tracing paper. Names, numbers or other symbols should also be written in India ink or, preferably, by 'Letra-set' or a similar system. Reduction of these figures should be taken into account when preparing the lettering. Titles and legends to the figures should be typed on a separate sheet. All figures should be numbered. Plates should be avoided, but when used, they should be numbered and supplied with informative title.

CORRESPONDENCE: on matters related to manuscripts should be addressed to the respective Editor or to the Secretary. Correspondence on manuscripts after acceptance or concerning page proofs should be addressed to the Secretary of the Editorial Board:

Prof. Pier Luigi Nimis, Dipartimento di Biologia,
cas. Università I - 34100 TRIESTE (ITALY)

ISSN 0394-9125

STUDIA GEOBOTANICA

An international journal

Vol. 11 1991

EDITORS

G. Estabrook - Ann Arbor, Mi
L. Iljanic - Zagreb
E. Mayer - Ljubljana
L. Orlóci - London, On.
F. Pedrotti - Camerino
S. Pignatti - Roma
A. Pirola - Pavia
J. Poelt - Graz
L. Poldini - Trieste
E. Wikus Pignatti - Trieste

EDITOR IN CHIEF

D. Lausi - Trieste

SECRETARY

P.L. Nimis - Trieste



**Dipartimento di Biologia
Sezione di Geobotanica ed Ecologia vegetale**

Università di Trieste

LICHENS AS BIOINDICATORS OF AIR POLLUTION BY SO₂ IN THE VENETO REGION (NE ITALY)*

P.L. NIMIS, A. & G. LAZZARIN, D. GASPARO

Keywords: Air pollution, Air quality, Bioindicators, Biomonitoring, Lichens, Italy, Veneto.

Abstract: This paper presents the results of a study on air pollution by SO₂ in the Veneto Region (north eastern Italy), using the "Calibrated Lichen Index of Air Quality" (I.A.P.) proposed by Liebendoerfer et al. (1988). This index, based on the frequency of epiphytic lichen species within a sampling grid of 10 units, has been tested by the authors in two preliminary studies carried out in two different areas of northern Italy (the town of La Spezia and the northern part of the province of Vicenza), revealing a high degree of correlation with pollution by SO₂, measured by automatic recording gauges. The index has been computed for 662 stations scattered throughout the Region of Veneto, whose surface area is of 18.364 km². The average number of relevés in each station is 3.7, with a total number of 2.425 relevés. The tree genus selected for sampling is *Tilia*. Altogether, 80 lichen species have been recorded during the survey, with a prevalence of nitrophytic and neutro-basiphytic lichens indicating a diffuse secondary eutrophication of the bark by agricultural activities.

Pollution maps for the seven provinces and for the entire region are presented. The evaluation of pollution levels is based on the correlations between the index and SO₂ levels found in the preliminary studies. All maps have been drawn by programs of automatic mapping.

The results show that in 93.7 % of the regional territory there is a very low probability of the occurrence of SO₂ concentrations higher than the thresholds established by law; this part of the region has yearly means of the 98th percentiles lower than 84 µg/m³. 25 % of the region has moderate pollution levels, 23.4 % a low pollution, and 44.9 % a very low air pollution. The most polluted areas are located in the southwestern part of the region. The pollution pattern agrees very well with the location of the main sources of pollution, and with the directions of the prevailing winds.

The study presents also the computerized distribution maps of eight selected species (*Candelaria concolor*, *Candelariella xanthostigma*, *Hypogymnia physodes*, *Parmelia exasperatula*, *Parmelia sulcata*, *Physcia adscendens*, *Physcia orbicularis*, *Xanthoria parietina*), that are discussed in relation with the main pollution patterns. The lichens species can be subdivided into three main groups, according to their distribution patterns. The main

* The publication of this study has been financed by a grant of the Consiglio Nazionale delle Ricerche to the first author.

factors affecting the distribution of lichens in the survey area are air pollution and the eutrophication of bark by agricultural activities.

A monitoring strategy where bioindicators and instrumental recording of air pollution are used jointly is proposed as a good solution to air pollution mapping over vast areas.

Riassunto breve: *LICHENI COME BIOINDICATORI DI INQUINAMENTO ATMOSFERICO NELLA REGIONE DEL VENETO (ITALIA NORD ORIENTALE).* - Questo lavoro presenta i risultati di uno studio sull'inquinamento da SO₂ nella Regione del Veneto, svolto utilizzando l' "Indice lichenico calibrato di qualità dell' Aria" (I.A.P.) proposto da Liebendoerfer et al. (1988). Tale indice, basato sulla somma delle frequenze delle specie licheniche epifite in un reticolo di 10 unità, è stato preliminarmente saggiato in due diverse aree dell' Italia settentrionale (città di La Spezia ed Alto Vicentino), rivelando un'alto grado di predittività rispetto alla contaminazione da anidride solforosa, misurata da apposite centraline. L'indice è stato calcolato per 662 stazioni (per un totale di 2425 rilievi) sparse per un'area di 18.364 km². Tutti i rilievi sono stati effettuati su Tiglio in un periodo di un anno, compreso tra il Settembre del 1989 ed il Settembre del 1990. Sono state rinvenute 80 specie di licheni, la maggior parte dei quali cresce di norma su substrati eutrofizzati, a reazione neutro-basica, ed è caratteristica dell'alleanza *Xanthorion parietinae*. Ciò indica una diffusa eutrofizzazione dei substrati dovuta in primo luogo all'intensa attività agricola.

Vengono presentate carte della contaminazione dell'aria da anidride solforosa per le sette provincie amministrative, e per l'intera Regione del Veneto. La stima dei livelli d'inquinamento si basa sulle correlazioni, ottenute negli studi preliminari, tra i valori di I.A.P. e le effettive concentrazioni di SO₂ in atmosfera rilevate da centraline. Le carte sono state ottenute attraverso programmi di cartografia automatica.

I risultati dimostrano che nel 93.7 % del territorio regionale la probabilità di superamento delle soglie di SO₂ stabilite per legge è bassa o molto bassa (medie annue dei 98^a percentili minori di 84 µg/m³). Il 25 % del territorio regionale presenta un moderato inquinamento atmosferico, il 23.4 % un inquinamento modesto, il 44.9 % ha livelli di inquinamento trascurabili. Le aree maggiormente inquinate, quelle cioè in cui maggiore è la probabilità del superamento delle soglie previste dalla legge, si trovano nella porzione sudoccidentale del Veneto, nelle provincie di Padova, Rovigo e Verona. L'andamento generale dell'inquinamento evidenziato dalle carte è in buon accordo con la localizzazione delle principali fonti inquinanti e con la direzione dei venti prevalenti.

Lo studio include anche le carte di distribuzione di otto specie licheniche (*Candelaria concolor*, *Candelariella xanthostigma*, *Hypogymnia physodes*, *Parmelia exasperatula*, *Parmelia sulcata*, *Physcia adscendens*, *Physcia orbicularis*, *Xanthoria parietina*), anch'esse ottenute per via automatica, che vengono discusse in relazione alla distribuzione spaziale dell'inquinamento atmosferico nel territorio regionale. Si possono distinguere tre gruppi principali di specie, con differenti patterns di distribuzione nell' ambito della regione. I fattori principali che condizionano la distribuzione dei licheni nella Regione Veneto sono l'inquinamento atmosferico e l'eutrofizzazione delle scorze conseguente all'attività agricola.

L'utilizzazione congiunta di bioindicatori e del rilevamento strumentale viene proposta come la strategia più appropriata al monitoraggio dell'inquinamento atmosferico su vasta scala.

INTRODUCTION

The aim of this paper is to present the results of a study on air pollution by sulphur dioxide in the Veneto Region (NE Italy) using epiphytic lichens as bioindicators.

This study, commissioned by the Veneto Region to the Department of Biology of the University of Trieste, is the first large-scale project on lichens as bioindicators in Italy, covering a surface area of 18.364 km². The field data were obtained in 662 sampling stations, for a total of 2.425 relevés of lichen vegetation, based on the method proposed by Liebendoerfer et al. (1988).

The survey, directed by the first author with the collaboration of Co.Ge.V. s.r.l. (Verona) and Ecothema s.r.l. (Trieste), has been completed within one year, from September 1989 to September 1990.

The maps presented in this paper can be used to detect high-risk areas, i.e. areas where the probability of concentrations of SO₂ higher than the thresholds established by law is greatest, and to optimize the positioning of the automatic recording gauges.

Air quality and air pollution

The terms "air quality" and "air pollution" have a different meaning. Air quality refers to the effects of different pollutants on a variety of subjects, including man, animals, plants and even inorganic substrates such as monuments, whereas air pollution is defined in terms of concentrations of pollutants in the atmosphere. As discussed by Nimis (1990), "air quality" is much more difficult to define on an operational basis than "air pollution", for four main reasons; these are:

- a) scarce knowledge on the effects of several pollutants on man, animals and plants,
- b) the fact that injury does not always depend on average pollution values, but also on peak levels and on the duration of exposure,
- c) scarce knowledge of the synergic effects of different pollutants on different organisms,
- d) scarce knowledge on the transfer of pollutants in the ecosystems.

The difficulties in establishing reliable air quality indices on a quantitative basis often led to search for "air quality indicators", which are parameters indicating the degree of air quality of a given site. In the absence of a clear operational definition of the term "air quality" the indicators become the only way to define it. This may involve circular reasoning.

Although the term "air pollution" is much easier to define on an operational basis, the monitoring of air pollution is far from being an easy task. The main reasons are:

- 1) The threshold levels established by law for the single pollutants are expressed as actual concentrations in the atmosphere. This requires a monitoring strategy based on direct instrumental recording.
- 2) The concentrations of pollutants in the air are highly variable in time and space. Monitoring must be based on a statistical approach, requiring a high density of sampling points both in time and space.
- 3) The high costs of instrumental monitoring are a strong constraint on the density of sampling points. Hence, data by instrumental nets often have a poor statistical quality, in spite of the apparent precision of the single measurements.

To overcome these problems, many different "pollution indicators" have been proposed. Their use is conceptually easier than that of air quality indicators, since air pollution is easily defined in a quantitative way. An indicator of air pollution is any kind of parameter which is known to be related to pollution levels. If this relation can be expressed by a highly significant mathematical relation the indicator can be used as a "monitor".

This study is based on lichens used as indicators. In the following we shall try to clarify what kind of information, relative to air quality and/or air pollution, can be obtained by our data.

Bioindicators of air pollution and air quality

The use of bioindicators of air pollution is based on the estimation of the effects of environmental changes due to air pollution on living organisms. An organism is considered as a bioindicator when it presents identifiable reactions to different degrees of pollution. It may react specifically to a certain pollutant or unspecifically to a mixture of toxicants. A bioindicator is optimal when it can be used as a biomonitor. Most studies dealing with bioindicators have a serious handicap: they do not provide quantitative pollution data since most bioindicators cannot be used as biomonitors. For this reason their use for regulatory decision-making may be problematic, since the maximum acceptable pollution levels are usually defined by law in terms of concentrations of the pollutants in the atmosphere.

Organisms may be also used as indicators of air quality. Their use is based on the assumption that the reaction of living organisms to pollution phenomena is a better estimate of damage to man or other animals than quantitative data on the concentrations of single pollutants. In this case the exact nature of pollution, the concentration of substances involved and the duration of pollution incidents are not determined, but real potentialities for damage will be identified. Although this is probably true in most cases, such an approach may

be subject to severe criticism: often there is no experimental evidence that the effects of pollution will have the same intensity on the bioindicator and on man, other animals, or other plants. When the use of a bioindicator of air quality is the only way to define air quality itself the approach, involving circular reasoning, cannot easily be accepted as scientifically valid.

Lichens as bioindicators - some problems

Epiphytic lichens are generally known as very good indicators of air pollution and/or air quality, and as such they have been extensively used throughout the world. For references on the very abundant literature on the subject we refer to Ferry et al. (1973), Lackovicova et al. (1988,1989), and Nimis et al. (1991). There are different techniques for the use of lichens as bioindicators; among the most widely used is the approach based on Indices of Atmospheric Purity (I.A.P.). Several different indices have been proposed; most of them associate a toxitolerance factor to each species. This factor is often calculated on a rather subjective basis, and it appears as one of the main weakpoints of several indices. When reliable, direct pollution data are not available, i.e. just in the case where a biomonitoring study is most useful, the attribution of toxitolerance values to each species may even involve circular reasoning; this happens when new sensitivity values towards pollution are used to obtain a number which is claimed to be related to pollution itself. Furthermore, lichens are very sensitive to climatic conditions, and in particular to air humidity. The "drought effect" (Beschel 1958, Rydzak 1969), i.e. the drier microclimate of large towns, may not be a severe limiting factor in regions with an oceanic-suboceanic climate, such as western Europe (see Barkman 1969, Coppins 1973), but its importance increases with increasing continentality of the general climate (Nimis, 1986). There is evidence that the tolerance of a given species to air pollution may differ according to the general climatic conditions. Thus, even in the rare cases where a toxitolerance factor for calculating an I.A.P. index is based on experimental data, its extension to another area, with different climatical conditions, is not always justified.

Recent studies in Switzerland by Herzig et al. (1985, 1987), Wanner et al. (1986), Liebendoerfer et al. (1988), Ammann et al. (1989), Urech et al. (1990) and Herzig (1990) led to the development of a new biomonitoring technique, based on epiphytic lichens, that proved to held very good results. These authors tested 20 different I.A.P. formulas against real pollution data. The sum of the frequencies of all species within a sampling grid of 10 subunits proved to have the highest statistical correlation with the combined concentrations of different pollutants. This simple index, called "Calibrated Lichen Index of Air Quality", has a great advantage over other I.A.P. indices, since it does not need the

attribution of toxitolerance values to each species. For these reason, this study is based on the I.A.P. index proposed by the above cited Swiss authors.

Predictive value of the adopted I.A.P. index

Liebendoerfer et al. (1988) tested the I.A.P. index later adopted in this study on the basis of direct pollution data derived from two independent measuring networks in Switzerland, one located in the town of Biel-Bienne, the other in the Swiss Mittelland. In Biel-Bienne these authors found a highly significant statistical correlation between the index and the sum of 8 different pollutants (dust, SO₂, NO₃, Cd, Cu, Pb, Cl, Zn). The predictability of the index decreased when some of the pollutants were excluded from the analysis. In the Swiss Mittelland the mixture of pollutants consisted of SO₂, NO, NO₂, O₃; also in this case the I.A.P. index proved to have a very high statistical correlation with the pollution data, and the significance of the correlations decreased when some pollutants were excluded from the analysis. The authors concluded that lichens are sensitive not only to SO₂, as it was generally assumed by most authors, but to a mixture of toxicants. As such, they can be utilized as monitors of air quality. In this case "air quality" is defined in terms of probability of recording concentrations of some pollutants that are above the threshold levels established by law. This definition does not involve circular reasoning, being based on a sound experimental design and on a statistical analysis of both lichen and pollution data.

Testing the method in Italy

Before adoption of the methodology proposed by the Swiss authors in the survey of the Region of Veneto, we decided to test it in a narrower area located within the north Italian territory. The area chosen was the town of La Spezia (Liguria) and its surroundings. The available pollution data were limited to SO₂, recorded by 8 gauges. Data on metal deposition have been obtained indirectly in 34 stations, using a lichen (*Parmelia caperata*) as a bioaccumulator. The results were as follows:

- 1) Regardless of the fact that in the immediate surroundings of the 8 gauges it was not possible to carry out all relevés in optimal standard conditions (absence of suitable trees), the I.A.P. showed a good statistical correlation with the SO₂ data (Nimis et al. 1990).
- 2) Some metals were highly correlated with the I.A.P. values, while others showed no statistically significant correlation. The correlation between I.A.P. values and the sum of SO₂ plus all metals was lower than that against SO₂ data alone (Nimis et al., unpubl.).

A still lower statistical correlation between I.A.P. and metal deposition patterns was found by us in a further test study, carried out in the northern part of the province of Vicenza, using lichens as bioindicators, and *Parmelia subrudecta* as a bioaccumulator of 10 metals (Nimis et al. 1991). This area is characterized by a very scattered occurrence of pollution sources within a mainly agricultural landscape: the use of lichens as bioaccumulators allowed the detection of the main sources for specific metals with a very high degree of accuracy, and the I.A.P. map was well in agreement with the location of the main sources of SO₂, and with the direction of the prevailing winds. However, there was a low correlation between I.A.P. index and the sum of the concentrations of all metals.

These results confirm the high predictive value of the Index with respect to SO₂, but do not agree completely with the model found by the Swiss authors. This might depend on two main reasons:

1) Different pollutants have different effects on lichen species; some species are able to tolerate high concentrations of some metals without apparent damage, but are much more sensitive to other gaseous pollutants.

2) When the emitting sources are concentrated in a relatively narrow space (e.g. an urban area and an adjacent industrial zone) the sum of their distribution patterns may reflect well the air quality situation, including possible synergic effects. On the other hand, when the emitting sources have a scattered distribution, some parts of the survey area may have high concentrations of a specific pollutant, while all other pollutants have low values. The effects of these concentrations on the I.A.P. values obviously depend mainly on the specific toxicity (for lichens) of the pollutant with the highest concentration.

In other words, it is more probable that the I.A.P. will show a highly significant correlation with the sum of several pollutants when their deposition patterns are similar. When this is not the case, such a situation cannot always be expected. Our knowledge on the toxicity of different pollutants for lichens is not yet sufficient to allow the assignment of a weight to every single pollutant, in order to evaluate its possible effect on the I.A.P. value.

In conclusion, the evidence which is available to date allows the use of the I.A.P. index adopted in this study as a reliable tool to estimate pollution by SO₂ and/or other phytotoxic gases. In the interpretation of our maps we have used the relation between I.A.P. index and SO₂ concentrations, as established at La Spezia (Nimis et al. 1990), to estimate the main pollution levels within the region. We think, however, that the index should be re-calibrated with pollution data relative to the Veneto Region, owing to climatical differences with La Spezia. Therefore, the translation of the index values into pollution values should be considered just as a first attempt to use lichens as biomonitors of SO₂ pollution in the region, and could be improved with further research. There are a few data available on SO₂ concentrations within the Veneto Region; these,

however, concern mostly areas located within the lichen desert, or with a very poorly developed lichen vegetation, and hence they do not allow a good calibration of the I.A.P. values. However, as far as the lowest I.A.P. values are concerned, there is a very good correspondence between the SO₂ concentrations recorded at La Spezia and those recorded within the Veneto Region.

It must be underlined that there is sufficient evidence to state that the index is linearly related to pollution by SO₂; the effect of a new calibration should be to modify slightly the slope of the regression line obtained at La Spezia. Considering that secondary eutrophication of the bark is much more pronounced throughout the Veneto Region than at La Spezia, and given that this factor may counterbalance the acidification of the bark due to air pollution, it is probable that a new calibration in the Region of Veneto will reveal somewhat higher pollution levels than those reported in this paper.

The maps which are presented here reflect the main distribution patterns of SO₂ pollution within the Region of Veneto.

Advantages of using bioindicators

One of the main advantages of the techniques based on bioindicators is their relatively low cost. This allows the adoption of sampling strategies with a relatively high density of sampling points, which, considering the nature of pollution phenomena, greatly enhances the data quality. The degree of error intrinsic in the variability of biological data is compensated by the density of sampling points. This permits the drawing of reliable pollution maps of large areas, that could not have been obtained by instrumental recording owing to the scarcity of sampling points.

However, the use of bioindicators should not always be considered as an alternative to instrumental recording. Its greatest advantages are evident when the two approaches are used jointly. One of the main problems of instrumental recording is the positioning of the recording instruments. Considering their generally low number, this is a most fundamental point, with a great influence on the results of the instrumental monitoring net. A large-scale survey of the area to be monitored by instrumental recording, using an appropriate bioindicator, can result in the relatively rapid production of a map showing high-risk areas, i.e. areas where the highest pollution levels are most probable. These areas can be selected for the positioning of recording gauges. In this way the use of monitoring instruments, which are necessary to measure the real extent of pollution levels, can be optimized.

An integrated approach, where bioindicators, bioaccumulators and recording instruments are used jointly, is probably the most appropriate solution for air pollution monitoring.

SURVEY AREA

The Region of Veneto, in north eastern Italy, has a surface area of 18.364 km² and a population of 4.380.252 inhabitants, with a density of 238 inhabitants/km².

The region is subdivided into 7 administrative provinces. The provincial capitals are: Belluno (population: 36.000), Padova (population: 222.150), Rovigo (population: 52.484), Treviso (population: 84.500), Venezia-Mestre (population 324.270), Verona (population: 258.700), and Vicenza (population: 109.500).

The northern part of the region is occupied mainly by mountains: the Dolomites in the province of Belluno, the Asiago Plateau in the province of Vicenza and the Lessinian Mountains in the province of Verona. The mountain area has the lowest population density: it hosts only 8.1% of the population, although it covers 29 % of the regional territory.

A low belt of hills connects the Alps and pre-Alps with the Venetian plain, which is interrupted by two other chains of hills of volcanic origin (Colli Euganei and Monti Berici); altogether the hills occupy 14.5 % of the regional territory and host 15 % of the population.

Most of the population (76.6 %) lives in the plain, where there is also the highest concentration of industries and an intensive agricultural activity; 56.4 % of the regional territory is located in lowland areas.


The main industrial areas are located in the lowlands, near Mestre-Marghera (mainly chemical industries), in the surroundings of Padova, Vicenza and Verona; large parts of the plains have a scattered, diffuse distribution of small industries, so that industrial and agricultural landscapes are often intermingled.

The main climatic features of the survey area are presented in fig. 1, that reports the climatic diagrams (according to Walter & Lieth 1960) of 17 selected localities, whose location is shown in fig. 2. Three main climatic types can be recognized:


1) Temperate climate (from Agordo to Cortina in fig. 1) : this climatic type is characteristic of mountain stations. Average yearly temperatures are lower than 13 °C and precipitation is the highest within the region, with yearly means always higher than 1000 mm. There is no sign of a decrease in water availability during summer months, precipitations being high from spring to autumn.

2) Intermediate climate (from Castelfranco to Vicenza in fig.1): this climatic type is clearly intermediate between types 1) and 3), and corresponds to the "esalpic climate" of Del Favero et al. (1990): it is characteristic of localities situated in the western and in the northern plains: there is a clear decrease of precipitation


AGORDO 9,3° 1366
(25-30) 611 m




ASIAGO 6,7° 1407
(30-34) 999 m




BASSANO DEL GRAPPA 12,7° 1177
(29-36) 129 m




BELLUNO 11,1° 1254
(30-30) 400 m




CORTINA D'AMPEZZO 6,5° 1159
(18-30) 1275 m




CASTELFRANCO VENETO 13,5° 915
(21-30) 44 m




PORTOGRUARO 12,6° 1121
(19-30) 6 m



TREVISO 13,4° 938
(30-30) 15 m



VICENZA 13,7° 1037
(30-30) 42 m






Fig. 1: climatic diagrams (following Walter & Lieth 1960) of 17 stations in the Region of Veneto; their locations are shown in fig.2. Further explanations in the main text.



Fig. 2: location of the meteorological stations whose climatic diagrams are shown in fig. 1, and directions of the winds in the Venetian Plain.

during summer, with two maxima in spring and autumn, but water availability is good throughout the year.

3) Mediterranean-like climate (from Badia Polesine to Verona in fig.1): it is characteristic of the southern plains. Average yearly temperatures are around 13 °C, and precipitation is generally lower than 800 mm. A brief period of summer drought is evident in all climatic diagrams, but a real water deficit does not occur. For this reason this climatic type cannot be considered as typically Mediterranean. The stations of Rovigo and Badia Polesine, the only ones where there is a short period of water deficit in summer, are located in areas with a very high water table. This explains the rarity in the region of true mediterranean vegetation, which is confined to coastal sand dunes. An important point, which is not visible in the climatic diagrams, is the stagnation of fog: this is a very frequent phenomenon during the colder seasons, particularly in the southern plains and in the province of Rovigo, and constitutes an important additional source of water for lichens.

Winds are an important factor in the dispersion of pollutants. Fig. 2 reports the main wind directions for 7 selected localities. The eastern plains are still under the influence of the Bora-winds, which are strong, dry, cold winds blowing from NE, particularly during the cold season. In the high plains (e.g. station of Bassano) local northern winds blowing from the pre-Alpine valleys can play a relevant role. In general, the direction and intensity of the winds in mountain areas are very variable, depending on the local morphology, and on the orientation of the main valleys. Near the coasts (e.g. station of S. Nicolò di Lido), there is a clear prevalence of southeastern winds coming from the Adriatic sea. The area around Padova is characterized by the prevalence of northern winds, whereas the three stations located in the western part of the region (Verona, Vicenza and Rovigo) have different wind patterns: Verona is characterized by the prevalence of eastern and western winds, Vicenza has a slight prevalence of northern winds, and Rovigo is still under the effect of the Bora winds from NE, which is much weaker than in the eastern stations. On the whole, the low plain is characterized by a counterclockwise rotation of the main wind directions.

DATA AND METHODS

Sampling

All relevés of lichen vegetation were on *Tilia* sp. This tree has been chosen for sampling since it is widespread throughout the region, both in natural stands (on the hills and lower mountains) and in urban areas.



Fig. 3: location of the 662 sampling stations within the regional territory.

Furthermore, the bark of *Tilia* has a subacid pH, ranging from 4.8 to 5.6, with an average of 5.2, and is of mesotrophic type (Barkman, 1958), which is ideal to estimate the compositional variations of epiphytic lichen vegetation due to acidification induced by SO₂.

The relevés have been carried out only in stations satisfying the following criteria:

- 1) At least 4 trees satisfying the standard features (see below) should be present.
- 2) The trees must have a circumference between 80 and 230 cm.
- 3) The trunks must not have an inclination higher than 10° (possibly below 3°).
- 4) The trees should be neither in closed natural stands, nor located adjacent to possible sources of disturbance, such as high-traffic roads, garbage cans, etc.

A total of 2.425 relevés was carried out in 662 stations, the location of which is shown in fig. 3.

The distribution of the stations in the 7 administrative provinces of the Region of Veneto is as follows:

Province of Belluno: 30 stations.

Province of Padova: 80 stations.

Province of Rovigo: 65 stations.

Province of Treviso: 80 stations.

Province of Venezia: 60 stations

Province of Verona: 120 stations.

Province of Vicenza: 227 stations.

The higher density of stations in the northern part of the province of Vicenza is due to the fact that in this area we carried out a preliminary survey to test the methodology which was later extended to the entire region, and in particular to estimate the optimal sampling density. The results of this survey, which show the degree of accuracy that can be reached by a higher sampling density, have been published by Nimis et al. (1991).

The sampling strategy was as follows:

- 1) The choice of the stations was based on preferential sampling because of the uneven distribution of suitable stations. A large amount of time was spent in the research of suitable stations, and in large parts of the survey area all stations fulfilling the standard features have been sampled.
- 2) In each station relevés were carried out on at least four randomly selected trees.
- 3) Relevés were taken using a sampling grid of 30 x 50 cm, subdivided into ten rectangles, placed on the boles at an height of 1.2 m. The centre of the grid was positioned in the part of the bole with highest lichen coverage.
- 4) A relevé listed all species found within the grid, with the number of grid units in which every species occurred (frequency value) being computed.
- 5) The sum of the frequency values of all species is the I.A.P. of the relevé.

6) When the I.A.P. values of the four relevés of a station were widely different, more relevés were carried out, until the distribution of the values fitted a bell-shaped curve. When no more trees were available, the station point was considered as a "low data-quality point".

7) The I.A.P. of the station is the average of the I.A.P. values of all relevés within the same station.

8) Low data-quality points were disregarded in the analysis unless no other suitable station was present within a radius of at least 10 km from the next high data-quality point. In any case, the number of low data-quality points does not exceed 5 % of the total number of stations.

Prior to sampling, the performance of the operators has been checked during a training period, so that the results obtained by different people were comparable.

A card has been compiled for each station, with the following information:

1) Geographical coordinates of the station and its location on a topographic map at a scale of 1: 25.000.000.

2) Schematic drawing of the sampling site with the exact location of the trees where relevés were carried out.

3) Colour picture of the sampling site.

4) Diameter and inclination of each tree, and exposure of the relevés on the trunks.

5) Total cover of lichen vegetation within the sampling area of each relevé.

The information in 1-3 will allow further repetitions to be made at a later date, to monitor the temporal variation of pollution phenomena throughout the region. These data are not reported here for reasons of space. They are available upon request to the Region of Veneto.

Data Analysis

In order to study the compositional variation of epiphytic lichen vegetation in the survey area the data have been processed by programs of multivariate analysis (classification and ordination). The analysis has been carried out in two steps:

1) For each station, the frequency data of each species in each sampling grid (relevé) have been averaged. A station vector is defined by the average frequency values of all species occurring in the relevés of the same station. The station vectors of each province have been separately subjected to numerical classification, in order to obtain groups of stations with similar floristic composition. The algorithm used was Complete Linkage Clustering with Euclidean Distance, using the program package of Wildi & Orloci (1983). These

results have been used to comment briefly on the main features of lichen vegetation within each province.

2) To obtain a general synthesis, a second matrix has been constructed, where the rows are species and the columns report the station groups obtained in the previous step; for each species the matrix reports its percentage occurrence within a station group. This matrix has been processed by numerical classification and reciprocal ordination, to detect the main trends of floristic variation within the survey area, and to discuss the main factors which are responsible of this variation. In this case also, the program package by Wildi & Orloci (1983) has been used.

The choice to subdivide the entire data set into 7 subsets (one for each province), and to use for the synthesis the contingency table of the groups obtained by numerical classification of the subsets, was based solely on the ground that it was necessary to reduce the computing load to perform multivariate analyses of the matrices.

Mapping procedures

Two types of maps are presented in this paper:

1) I.A.P. maps, showing the spatial distribution of the I.A.P. values within each of the seven provinces, and within the entire Veneto Region. These are based on the I.A.P. values of the different stations.

2) Maps showing the distribution patterns of selected species. These are based on the average frequencies of each species within each station.

All maps have been obtained by computerized automatic mapping in order to avoid subjectivity in the cartographic expression of the results. We have utilized the program package SURFER (Golden Software Inc., Golden, Colorado, U.S.A.). The programs are based on a grid, whose mesh size depends on the minimum and maximum values of the input data. The interpolation method utilized to create a regular grid starting from points (stations) irregularly distributed on the survey area is based on the I.A.P. values of the 10 stations nearest a given point. The influence of a given station on the others is inversely proportional to their distances.

In interpreting the maps, one should be aware that their reliability is not the same for each part of the map, since it depends on the density of sampling points. The density should be higher in mountain areas with a rugged morphology. However, it is just the mountain area that has been sampled with lowest intensity in our study. The main reason for this choice is that we have preferred to attain a higher sampling intensity in the more densely populated areas of the Venetian Plain. The maps of the mountain areas, and in particular that of the province of Belluno, are mostly valid only for sites located in the main valleys. No station was located above 1.300 m.

Taxonomical remarks

Since all relevés have been carried out in the field, some species presented identification problems that cannot always be solved without analysis in the laboratory. In most cases, dubious material was collected in the field, and identified later in the laboratory. In the following we list a few cases in which the adopted nomenclature, which mostly follows Nimis & Poelt (1987) needs clarification.

- *Buellia punctata*: the epiphytic forms of the *B. punctata* complex are still very poorly known, and in need of revision. Here the species is meant in a very broad sense.

- *Lecanora* gr. *hagenii*: this name includes two lichens, *L. hagenii* s.str., which occurs mainly on eutrophic bark in relatively unpolluted areas, and another lichen, belonging to the same group, which, however, differs from *L. hagenii* s. str. in the non-pruinose apothecia with entire margin. The latter is most common in polluted areas, and sometimes is the only species which occurs within urban settlements; perhaps it corresponds to forms which are sometimes treated as *Lecanora umbrina* auct.

- *Lecidella elaeochroma*: this species belongs to a difficult group, where several species have been recognized on the basis of characters whose taxonomic significance is still not clear; in the survey area *Lecidella elaeochroma* s.str. is the most frequent taxon, but also *Lecidella achristotera* and *Lecidella euphorea* have been recorded sometimes. Since these three species are difficult to distinguish in the field, they have not been distinguished in the relevés, and here *L. elaeochroma* is meant sensu lato.

- *Nectriacea* sp.: with this name we designate a conidiophoral lichenized fungus with small bright yellow pycnidia and a poorly developed greenish thallus, which appears to be relatively frequent in some parts of the region, even in rather polluted areas. The identification at genus and species levels has not yet been possible (Poelt, in litt.).

- *Physcia vitii*: with this name we designate a *Physcia* resembling *Physcia adscendens*, but with a more robust thallus lacking fibrils. It corresponds well with the description of *P. vitii*, but we have no reference material to check this identification.

- *Rinodina exigua*: this is the most common epiphytic *Rinodina* throughout the survey area. However, some samples identified in the laboratory proved to belong to *R. pyrina*. Since the two species are difficult to distinguish without the observation of the spores, some citations of *R. exigua* might actually refer to *R. exigua*.

- *Usnea subfloridana* aggr.: the thalli of *Usnea* found throughout the survey area are always very poorly developed; this identification cannot be other than an "educated guess" and it might be that more taxa are involved here.
- *Xanthoria fallax*: according to J. Poelt (in litt.), who is preparing a revision of *Xanthoria fallax* s. lat., the material identified as *X. fallax* by European lichenologists includes at least three distinct species: of these, *Xanthoria fallax* s. str. *Xanthoria fulva*, and another taxon provisionally named *Xanthoria bella* occur in our survey area; they have not been distinguished in the relevés, so that here *Xanthoria fallax* is treated in the old sense, including the three species.

RESULTS

The results of this study are presented in this chapter, according to the following scheme:

- 1) Numerical analysis (classification and ordinations) of the matrix of species and station vectors, relative to all provinces.
- 1) For each of the seven provinces: (a) classification of the station vectors into groups, and (b) I.A.P. maps of the provinces.
- 3) I.A.P. map of the entire region.
- 4) Distribution maps of selected species.

The basic data (2.425 relevés and the exact location of each station) are not presented in this paper for reasons of space. They are available upon request to the first author.

The lichen vegetation within the region

Altogether, 80 species have been recorded during the survey. The most common are, in order of decreasing frequency (see tab. 1): *Physcia orbicularis*, *Physcia tenella*, *Lecidella elaeochroma*, *Physcia adscendens*, *Xanthoria parietina*, *Lecanora chlorotera*, *Parmelia exasperatula*; all them have their optimum within communities of the *Xanthorion parietinae*, which, considering the physico-chemical properties of *Tilia* bark, indicates a strong secondary eutrophication of the substrates throughout the region. This remark concerns the whole flora, since most of the rare species are also *Xanthorion* elements: the eutrophication of the bark is mainly due to the intense agricultural activity.

The frequency vectors of all groups of stations obtained by the numerical classification of the 7 stations sets (one for each province, tab. 1) have been submitted to classification and ordination, in order to:

- 1) obtain a synthetic view on the lichen vegetation on *Tilia* in the survey area;
- 2) detect the main trends of compositional variation, and analyze the main factors underlying this variation.

Group n°	11122223333333333333333344444444444444
Province	VVVVVVTTTTBBBVVVVVVVVVVVVPPPPRRRRR RRRRRVVVVVLLLLIIIIIIIEEEEDDDDDOOOOO
Station group n°	123451234512312345671234123412345
Candelariella xanthostigma	233 33 11 1 1111
Physcia stellaris	2441 33 1 11 11
Xanthoria fallax	2322122 11 1 1 1 11
Parmelia subargentifera	231 12 1 1
Rinodina exigua	112122 1 1 1 11112 11
Chaenotheca furfuracea	121333 1 2 111 12 1
Physcia aipolia	4541222 1 1 11111 1
Lecanora carpinea	4551114 1 1111 1 111111 11 1
Lecanora gr. hagenii	3341 2 111111 21112221 11
Caloplaca cerina	2351 1 1 1 11 1
Usnea subfloridana	21133511 111 12 1
Evernia prunastri	12334111212221 13 1 1 111 1
Hypogymnia physodes	1221354212111111 12 11121121
Physcia vitii	11 12 3 1111 2111212111114
Parmelia subrudecta	12 23 11 211111 3 31 11231112 1
Lecanora subfuscata	23 1 2 111
Lecanora allophana	33 1 1 11 1 1 11 1
Candelariella vitellina	34 1 22
Parmelia tiliacea	14 11 31 1 1 11111 1 1
Parmelia pastillifera	12
Caloplaca ferruginea	12
Parmelia quercina	12 1
Parmelia glabra	21 1
Caloplaca flavorubescens	112 1 1
Parmelia saxatilis	1
Anaptychia ciliaris	1
Cliostomum corrugatum	1 1
Ramalina farinacea	11 1
Normandina pulchella	1 111
Physconia deterosa	1 1
Collema nigrescens	1
Physconia perisidiosa	1
Parmelia glabratula	1
Hypogymnia bitteriana	1
Physconia distorta	11
Opegrapha lichenoides	1
Arthopyrenia lapponina	1
Physcia clementei	1
Scoliciosporum umbrinum	1
Pertusaria amara	1

Heterodermia speciosa	1																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																				
-----------------------	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

Tab.1: Station vectors obtained by numerical classification of the 7 matrices of stations and species (one for each province), ordered as in the dendrogram of fig. 4. The numbers refer to 5 frequency classes, with intervals of 20 % each. Four groups are distinguished, as in fig.4. Provinces: BL (Belluno), PD (Padova), RO (Rovigo), TV (Treviso), VE (Venezia), VI (Vicenza), VR (Verona). Further explanations in text.




Fig. 4: dendrogram obtained by numerical classification of the matrix of station groups and species (tab.1). The station groups have been obtained by numerical classification of 7 matrices of stations and species, one for each province. The abbreviations of the provinces are as in the caption of tab.1. Further explanations in the main text.

The dendrogram of the frequency vectors is shown in fig. 4: four main groups are formed (1,2,3,4); tab 1 reports the species and the frequency vectors, ordered as in the respective dendrograms.

In the following, each group will be briefly discussed, referring also to the pollution pattern shown by the I.A.P. map (fig. 21), that will be commented on later.

Group 1: it includes only three groups of stations, located on the low hills of the provinces of Verona and Vicenza. In this group the number of high-frequency species is great. Strongly acidophytic species are less frequent than in the following group, whereas nitro- and neutro-basiphytic species, such as *Candelariella reflexa*, *Xanthoria parietina* and *Xanthoria fallax* are more frequent. Also in this case *Xanthorion* species are prevalent: some of them, however, are characteristic of moderately eutrophicated, subneutral-subacid bark (e.g. *Physcia stellaris*, *Physconia distorta*, *Parmelia subargentifera*). Also some species of *Parmelion* occur in this group, that represents well developed stands in relatively pollution-free areas with a moderate impact of agricultural activity (slight eutrophication of the bark).

Group 2: it includes four groups of stations, located in relatively undisturbed sites of the provinces of Belluno and Vicenza, mostly in the mountains. It has a large number of high-frequency species, and the greatest frequency of acidophytic and relatively anitrophytic lichens, such as *Chaenotheca furfuracea*, *Usnea subfloridana* aggr., *Evernia prunastri*, *Hypogymnia physodes*, *Parmelia sulcata*. The majority of the most frequent species, however, are elements of *Xanthorion* vegetation. This group represents well developed stands in relatively pollution-free areas where the impact of agriculture, and hence the secondary eutrophication of the bark, are low.

Group 3: this group includes 12 frequency vectors, relative to stations located in the provinces of Treviso, Venezia, Padova, Verona and Rovigo. Most of the stations are located in heavily urbanized areas, often close to industrial centres. This group is characterized by a low number of species, the most frequent ones being *Physcia orbicularis*, *Physcia tenella*, and *Lecidella elaeochroma*. Furthermore, there is a sporadic occurrence of acidophytic species such as *Hypogymnia physodes*, *Evernia prunastri*, *Usnea subfloridana*, *Lecanora conizaeoides*, *Parmelia exasperatula*, that are absent or much less frequent in the following group. This group represents a very impoverished epiphytic vegetation in urban or industrial areas where the impact of air pollution is high. The low frequency of highly nitrophytic species, such as *Candelariella reflexa*, *Hyperphyscia adglutinata*, *Candelaria concolor*, and the presence of a set of acidophytic species, indicates a lower impact of agricultural activities and a secondary acidification of the bark by air pollution.

Group 4: this group includes 14 frequency vectors, relative to stations located in all provinces except Belluno and Rovigo. Its floristic composition very




Fig. 5: Reciprocal ordination of the species (a) and of the station groups (b) based on the data in tab.1. In Fig. 5a the species with low scores on either axis are not discriminated, in fig. 5b the station groups obtained by numerical classification are designated by numbers, as in fig.4.

much resembles that of group 1, with the difference that most of the species have lower frequency values. This group represents relatively disturbed *Xanthorion* stands in areas with intense agricultural activity and a moderate air pollution.

The reciprocal ordination of frequency vectors and species, based on the data of tab. 1 is shown in fig. 5a (vectors), and 5b (species). In fig. 5a the clusters of vectors obtained by numerical classification mainly occupy four distinct quadrants in the space defined by the first two canonical variates. The first variate can be interpreted as a diversity axis, with species-rich vectors having positive scores, and species-poor vectors negative scores. The second variate reflects the degree of eutrophication of the substrates: groups 1 and 4, characterized by a high frequency of neutro-basiphytic and nitrophytic species, have positive scores, whereas groups 2 and 3, with a higher frequency of acidophytic and anitrophytic species, have negative scores. This interpretation is confirmed by the ordination of species, which is shown in fig. 5b: the species with low scores on both axes are either low-frequency species, or species not significantly associated with the main floristic variation trends, and are not discriminated in the analysis. The sequence of the indicator species (i.e. those with high scores on at least one canonical variate) along the first axis reflects a gradient of sensitivity to air pollution, the most resistant species being *Physcia tenella* and *Physcia orbicularis* (highest negative scores). At the positive and negative extremes of the second axis are, respectively, three highly nitrophytic species (*Candelaria concolor*, *Hyperphyscia adglutinata* and *Candelariella reflexa*), and a set of anitrophytic and acidophytic species (*Evernia prunastri*, *Parmelia exasperatula*, *Hypogymnia physodes*, *Usnea subfloridana* aggr.).

The reciprocal ordination of frequency vectors and species has been carried out also on a reduced matrix, without the groups of vectors 1 and 2, in order to analyze in more detail the floristical variation trends relative to the stations located in more polluted areas (groups 3 and 4). The results are shown in fig. 6a (groups) and 6b (species). In fig. 6a the groups 4 and 3 are well separated on the first canonical variate, the former having mostly positive scores, the latter negative scores. The group 3 has a narrower spread on the second variate, which is mainly due to the lower number of species, and hence to its higher within-group homogeneity. The arrangement of the indicator species in the space defined by the first two variates is shown in Fig. 6b: it is evident that the first canonical variate reflects a gradient of increasing acidity of the substrate, from positive to negative scores: acidophytic species, such as *Evernia prunastri*, *Parmelia sulcata*, *Parmelia exasperatula*, *Hypogymnia physodes*, *Usnea subfloridana*, *Chaenotheca furfuracea*, *Lecanora conizaeoides*, have negative scores on the first axis. The two pollution-resistant species evident in the ordination based on the entire data set, i.e. *Physcia tenella* and *Physcia orbicularis*, have a slightly different position in the ordination of fig. 6b: *Physcia tenella*




Fig. 6: Reciprocal ordination of the species (a) and of the station groups (b) based on the data in tab.1, without the groups of stations 1 and 2. In Fig. 6a the species with low scores on either axis are not discriminated.

occupies the negative extreme of the first axis, showing a higher resistance to acidification than *P. orbicularis*. The first canonical variate in the ordinations of Fig 6 shows a gradient of acidification, paralleled by a progressive decrease both in the number of species and in their overall frequency.

These results can be briefly summarized as follows: the most logical sequence of the groups of frequency vectors is: 2,1,4,3, because this sequence shows the main floristic variation trends along a gradient of increasing disturbance:

- Group 2 represents a situation where neither the pollution by phytotoxic gases nor that by organic and inorganic fertilizers have a great impact on the epiphytic lichen vegetation. This is shown by the relatively great number of high-frequency species (indicating low pollution levels), and by the presence of a set of anitrophytic and acidophytic species (indicating low eutrophication of the bark).

- Group 1 represents an intermediate situation, where air pollution by phytotoxic substances is still low (great number of high-frequency species), but there are slight signs of bark eutrophication by fertilizers (absence of anitrophytic species).

- Group 4 represents a situation where both the pollution by phytotoxic substances and by fertilizers have clear effects on the lichen vegetation: the number of high-frequency species is strongly reduced, and there is a maximum of nitrophytic lichens.

- Group 3 represents the final stage of the process, preluding to the lichen desert: the pollution by phytotoxic substances (chiefly SO₂) is so strong that, besides a further drastic reduction of high-frequency species, there is also a secondary acidification of the bark; this creates conditions which reduce the presence of neutro- basiphytic and nitrophytic species and favour the presence of some pollution-tolerant acidophytic species which occur also in group 2.

This scheme seems to be a general rule in the north eastern Italian Plains. Nimis (1986) in his study of the epiphytic lichen vegetation in the town of Udine (high Friulian Plain) described a situation very similar to that discussed above. In Udine a narrow belt of scarcely developed acidophytic species is present all around the lichen desert occupying the town centre. These species are otherwise absent or very rare in the Friulian Plain because of the intense agricultural activity, and the consequent pollution by fertilizers. They start to be common again only in hill and mountain areas where the agricultural activity is less intense. The disjunct distribution of *Pseudevernia furfuracea* in the northern part of the province of Vicenza (low mountains and main urban settlements), as shown by the map published by Nimis et al. (1991), is another good example of the secondary habitats created by air pollution for acidophytic species.

Province of Belluno

The province of Belluno occupies the northernmost portion of the region, and has a surface area of 3.678 km², being the largest province in the region. The population is of 215.000 inhabitants with a density of only 58 inhabitants/km². Most of the provincial territory lies in mountain areas, and the main urban centres are in the valleys. Industrial activities are scarcely developed, and the main industries are located in the surroundings of the provincial capital, Belluno, a town of 36.000 inhabitants.

The province of Belluno has the highest species richness (54 species). The most common species are *Lecidella elaeochroma*, *Parmelia sulcata*, *Parmelia exasperatula*, *Physcia tenella*, and *Evernia prunastri*. The classification of the 30 stations produced 3 main groups (tab. 2): acidophytic species, such as *Hypogymnia physodes*, *Parmelia sulcata*, *Parmelia exasperatula*, *Chaenotheca furfuracea*, *Evernia prunastri* tend to decrease from group 1 to group 3, whereas in the latter there is an higher frequency of neutro-basiphytic and nitrophytic species such as *Candelaria concolor*, *Physcia adscendens*, *Hyperphyscia adglutinata*. The sequence from group 1 to group 3 reflects a gradient of increasing eutrophication of the bark. In this province acidophytic species have the highest frequency, which is probably due to the fact that here there is the lowest agricultural activity in the region.

This is the province where our air quality map has the lowest reliability: the density of sampling stations is very low (see figs. 3 and 7), and it appears to be still more unreliable considering that most of the provincial territory is occupied by high mountains (all relevés have been carried out in the more densely populated valleys). The low density of sampling stations is due to two main reasons: a) the difficulty of finding suitable trees (*Tilia* becomes rare in mountain areas) and, b) the fact that most of the province has a low population density and a low concentration of industrial activities; we have preferred to give a higher sampling intensity to the densely populated high-risk areas of the plains.

With these limitations in mind, the map of fig.8 can be briefly commented on as follows: the province of Belluno has the highest I.A.P. values within the region, most of them being higher than 30; no area with lichen desert has been detected within the Provincial territory. The most polluted areas are in the southern part of the province, south of Belluno, and most of the pollution is likely to derive from Belluno itself and from its small industrial area. The area with lowest I.A.P. values (lower than 25) has an elliptical shape, with the longer axis oriented in a N-S direction, which accords well with the direction of the prevailing winds in the high Piave valley. A study on lichens as bioindicators in the high Piave Valley has been published by Caniglia et al. (1978) and Spampani (1982).

Station group n°	1	2	3
Lecanora gr. hagenii	2		1
Caloplaca holocarpa	1		
Xanthoria parietina	2	1	1
Lecidella elaeochroma	5	4	4
Parmelia sulcata	5	5	3
Parmelia exasperatula	5	5	3
Hypogymnia physodes	5	3	1
Physcia stellaris	3		1
Lecanora chlarotera	3	3	3
Physcia tenella	3	4	2
Rinodina exigua	2	2	1
Candelariella reflexa			2
Parmelia exasperata			2
Chaenotheca furfuracea	3	3	1
Physcia vitii	1		
Parmelia subaurifera	2		
Caloplaca cerina	1		1
Lecanora carpinea	1	1	1
Lecanora subfuscata	2		1
Usnea subfloridana	3	3	1
Parmelia tiliacea		1	1
Lecanora symmicta	2		
Arthopyrenia lapponina	1		
Evernia prunastri	3	3	2
Physconia distorta	3		
Parmelia subargentifera	1		
Pseudevernia furfuracea			1
Lecanora conizaeoides			1
Opegrapha lichenoides	1		
Graphis scripta	1		
Parmelia caperata		1	1
Normandina pulchella			1
Ramalina farinacea		1	1
Parmeliopsis ambigua		1	
Scoliciosporum umbrinum			1
Lecanora allophana	1		1
Opegrapha atra		1	
Nectriacea sp.		1	1
Arthopyrenia antecellens			1
Physcia biziana		1	1
Physcia aipolia	2	2	1
Arthopyrenia punctiformis	2		1
Parmelia subrudecta		3	2
Xanthoria fallax	2	1	2
Physcia hirsuta			1
Hyperphyscia adglutinata	1	2	4
Physconia grisea		2	1
Arthonia radiata	2	3	3
Candelariella xanthostigma	3	2	4
Physcia orbicularis	5	5	5
Candelaria concolor	3	5	5
Physcia adscendens	4	5	5

Tab.2: Frequency vectors of the three station groups obtained by numerical classification of the matrix of species and stations relative to the province of Belluno. Numbers refer to 5 frequency classes of 20 % each.




Fig. 7: map of the province of Belluno, with the location of the sampling stations.




Fig. 8: I.A.P. map of the province of Belluno. The province has been subdivided into several zones, characterized by different I.A.P. values.

Province of Treviso

The province of Treviso has a surface area of 2.477 km² and a population of 736.400 inhabitants, with a density of 297 inhabitants/km². Most of the territory (63.6 %) is located in lowland areas, with the exception of its northwestern part (29.4 %), which is occupied by low hills and mountains, the latter located only at the border with the province of Belluno. The main industrial area lies around the provincial capital, Treviso, a town of 84.500 inhabitants. Minor industrial areas are located near medium-sized urban centres, such as Conegliano, Montebelluna, Valdobbiadene, Vittorio Veneto (see fig. 9).

This is a rather species-poor province, with only 39 species recorded. The most frequent are *Physcia orbicularis* (95 % of the stations), *Candelaria concolor* (82 %), *Physcia tenella* (76%), *Xanthoria parietina* (60 %), *Lecanora chlarotera* (52 %); all of them are linked to *Xanthorion* vegetation. The numerical classification of the 80 stations produced 5 main groups that are shown in tab. 3. The groups of stations 1 and 2 are very species-poor; group 2 differs in the higher frequency of some acidophytic species such as *Parmelia sulcata*, *Hypogymnia physodes*, *Chaenotheca furfuracea*, *Usnea subfloridana*, *Evernia prunastri* and *Parmelia exasperatula*; both groups include stations located near industrial areas or within urban settlements, and represent, respectively, impoverished *Xanthorion* stands, and a species-poor vegetation characterized by the occurrence of acidophytic species due to secondary acidification of the bark. The remaining three groups of stations have a very similar floristic composition, and can be considered as representing three facies of *Xanthorion* vegetation in relatively unpolluted areas.

The I.A.P. map (fig. 10) shows that most of the province has relatively high I.A.P. values (25-50) indicating moderate, or low degrees of air pollution. An exception is the area south of Treviso, the largest urban and industrial agglomeration of the province, which includes a narrow lichen desert surrounded by areas with I.A.P. values lower than 20. Furthermore, minor areas with low I.A.P. values are evident southwest of Conegliano and Castelfranco Veneto, near Miane, Volpago and S. Biagio (see fig. 9); most of them include industrial agglomerations. It is rather surprising that the surroundings of Vittorio Veneto, a relatively large settlement with an industrial area, show the highest I.A.P. values in the whole province (higher than 40): this might be due to the local microclimate, since Vittorio Veneto is located at the end of a narrow valley with frequent N-S winds, which might help the dispersion of pollutants toward the south. The influence of Vittorio Veneto could be evident in the northward extension of the relatively polluted area surrounding Conegliano (I.A.P.: 21-25), which otherwise would be difficult to explain considering the main wind directions.

Station group n°	1	2	3	4	5
Physcia orbicularis	5	3	5	5	5
Candelaria concolor	3		5	5	5
Xanthoria parietina	3	2	4	4	4
Lecanora chlarotera	3	2	4	4	4
Parmelia sulcata	1	5	2	2	1
Hypogymnia physodes		2	1	1	1
Chaenotheca furfuracea	1	2	1	1	
Lecanora gr. hagenii	1		2	1	1
Caloplaca cerina			1	1	1
Physcia aipolia			1		1
Lecanora carpinea			1	1	1
Usnea subfloridana		2			
Evernia prunastri		3		1	1
Xanthoria fallax					1
Physconia grisea				1	1
Physcia stellaris			1		
Parmelia exasperata	1				
Parmelia caperata				1	
Lecanora subfuscata			1		
Catillaria nigroclavata				1	
Parmelia tiliacea			1	1	
Nectriacea sp.			2		
Parmelia acetabulum				1	
Arthopyrenia antecellens	1				1
Rinodina exigua	1	1	1		
Physcia biziana	1	1	1	1	1
Physcia hirsuta					1
Arthopyrenia punctiformis			1	1	1
Candelariella xanthostigma	1		2	3	1
Parmelia exasperatula	1	4	4	3	1
Candelariella reflexa			3	2	1
Physcia vitii	1		1	1	1
Parmelia subrudecta			2	1	2
Normandina pulchella	1				1
Arthonia radiata	2		2	3	4
Lecidella elaeochroma	3	3	4	3	2
Hyperphyscia adglutinata	1		2	3	4
Physcia adscendens	1		5	3	2
Physcia tenella	4	5	5	5	3

Tab.3: Frequency vectors of the five station groups obtained by numerical classification of the matrix of species and stations relative to the province of Treviso. Numbers refer to 5 frequency classes, with intervals of 20 % each.




Fig. 10: I.A.P. map of the province of Treviso. The province has been subdivided into several zones, characterized by different I.A.P. values.

Province of Venezia

The province of Venezia occupies most of the coastal part of the region; it has an area of 2.460 km², and a population of 833.600 inhabitants, with a density of 338 inhabitants/km². All of the provincial territory is below 100 m. In the Mestre-Marghera area, close to Venice, there is one of the largest industrial agglomerations of Italy, with a prevalence of chemical industry. Another industrial area, of much smaller size, is located near Portogruaro (see fig. 11).

In this province only 34 species have been recorded; the number of sporadic species, however, is rather low in comparison with other species-poor provinces. The most frequent species are: *Physcia orbicularis* (80 % of the stations), *Physcia tenella* (75 %), *Lecidella elaeochroma* (67 %), *Candelaria concolor* (53 %), *Lecanora carpinea* (50 %). All of them are more or less linked to *Xanthorion* vegetation. The numerical classification of the 56 stations produced 4 main groups, which are reported in tab. 4. The groups of stations 1 and 2 are very similar, and are characterized by the high frequency of *Lecanora carpinea*, *Candelaria concolor*, *Physcia adscendens*, *Hyperphyscia adglutinata*, *Arthonia radiata* and *Xanthoria parietina*: they represent relatively well developed *Xanthorion* stands; group 1 differs from group 2 in the higher frequency of *Parmelias*, such as *P. caperata*, *P. sulcata*, *P. subrudecta*, *P. exasperatula*. The groups of stations 3 and 4 have no clear differential species: they are characterized by having, in general, lower frequencies for most species, and represent impoverished *Xanthorion* stands; group 4 differs from group 3 in the higher frequency of some acidophytic species, such as *Parmelia sulcata*, *Hypogymnia physodes*, *Parmelia exasperatula*, and represent an impoverished *Xanthorion* vegetation in which secondary acidification by air pollution starts to become evident (most of these stations are located near industrial areas or large urban settlements).

The I.A.P. map (fig. 12) shows a marked difference between the eastern and western parts of the province. The eastern part is characterized by relatively high I.A.P. values (mostly between 30 and 50) indicating a moderate degree of air pollution; an exception is given by an area south of Portogruaro, corresponding to an industrial agglomeration, where there is even a small lichen desert, and which extends its influence to the easternmost part of the province. On the contrary, in the western part of the province the I.A.P. values are mostly lower than 20. An extended lichen desert occurs southwest of the huge industrial agglomeration of Marghera, which is certainly the most important pollution source. Its effects extend to the Lido of Venice, and most probably also to the provinces of Padova and even Rovigo. The only areas which are relatively less polluted are the surroundings of Chioggia, in the southern part of the province, along the coasts.

Station group n°	1	2	3	4
<i>Lecanora carpinea</i>	5	5	2	1
<i>Candelaria concolor</i>	5	5	2	1
<i>Physcia adscendens</i>	5	4	2	2
<i>Parmelia sulcata</i>	3	2	2	4
<i>Hypogymnia physodes</i>			1	2
<i>Physcia biziana</i>	1		1	
<i>Parmelia caperata</i>	2			2
<i>Cliostomum corrugatum</i>		1		
<i>Buellia punctata</i>		1		
<i>Catillaria nigroclavata</i>		1		
<i>Usnea subfloridana</i>			1	
<i>Physcia clementei</i>			1	
<i>Rinodina exigua</i>		1	1	
<i>Parmelia subrudecta</i>	3	1	1	
<i>Lecanora gr. hagenii</i>			1	
<i>Evernia prunastri</i>		1	2	1
<i>Arthopyrenia antecellens</i>		1	1	1
<i>Chaenotheca furfuracea</i>			1	
<i>Physconia grisea</i>		1	1	
<i>Opegrapha atra</i>			1	
<i>Lecanora allophana</i>	1	1	1	
<i>Lecanora carpinea</i>	1	1	1	
<i>Candelariella reflexa</i>	2	1	1	1
<i>Physcia stellaris</i>				1
<i>Parmelia tiliacea</i>		1	1	1
<i>Parmelia exasperatula</i>	4	1	2	4
<i>Candelariella xanthostigma</i>	5	1	1	
<i>Hyperphyscia adglutinata</i>	5	3	1	
<i>Arthonia radiata</i>	5	4	1	1
<i>Xanthoria parietina</i>	5	5	2	2
<i>Physcia vitii</i>	2	4	1	3
<i>Lecidella elaeochroma</i>	4	4	4	4
<i>Physcia tenella</i>	5	2	5	1
<i>Physcia orbicularis</i>	5	5	4	3

Tab. 4: Frequency vectors of the four station groups obtained by numerical classification of the matrix of species and stations relative to the province of Venezia. Numbers refer to 5 frequency classes, with intervals of 20 % each.




Fig. 12: I.A.P. map of the province of Venezia. The province has been subdivided into several zones, characterized by different I.A.P. values.

Province of Padova

The province of Padova has a surface area of 2.142 km², and a population of 818.650 inhabitants, with a density of 382 inhabitants/km², the highest within the region.

Most of the provincial territory (94.2 %) is below 100 m, with the exception of its southwestern part, west of Monselice (5.8 %), that is occupied by a chain of low hills (Colli Euganei). There is intense agricultural activity throughout the province, while most of the industries are concentrated around the city of Padova and along the highways connecting Padova with Monselice (see fig. 13).

This is a very species-poor province, with 27 species, most of which have been recorded only within a few of the 75 stations. The most frequent species are *Physcia tenella*, *Physcia orbicularis* and *Lecidella elaeochroma*, which occur in more than 50 % of the stations, followed by other *Xanthorion* species such as *Candelaria concolor*, *Xanthoria parietina*, *Candelariella reflexa* and *Lecanora chlorotera*, occurring in more than 20 % of the stations.

The numerical classification of the stations produced 4 main groups, reported in tab. 5. The groups have a very similar floristic composition, differing chiefly in the presence of a set of acidophytic species such as *Parmelia sulcata*, *Evernia prunastri*, *Hypogymnia physodes*, *Parmelia exasperatula*, *Pseudevernia furfuracea*, *Usnea subfloridana*, and *Lecanora conizaeoides*, their frequencies decreasing from group 1 to group 4, whereas nitrophytic species such as *Physcia orbicularis*, *Candelaria concolor*, *Xanthoria parietina*, *Candelariella reflexa*, *Hyperphyscia adglutinata* and *Physcia adscendens* have an opposite behavior. This may reflect the opposing effects on lichen vegetation of the secondary acidification of the bark in heavily polluted areas, and of the eutrophication due to agricultural activity.

The I.A.P. map of the province is shown in fig. 14. In general, the I.A.P. values are low, which indicates a moderate to high air pollution. The northern part of the province, and part of the Colli Euganei east of Monselice, are the only areas with I.A.P. values higher than 20. This is well in accordance with the direction of the winds which prevail over the city of Padova; they are mostly coming from the north (see fig. 2), and transport southwards the pollution produced by Padova and its industrial area. The small areas with I.A.P. values lower than 20, located in the northern part of the province, reflect the influence of the neighbouring towns of Bassano (Vicenza) and Castelfranco Veneto (Treviso) and of their industrial areas, both located north of the province of Padova.

The effects of the large industrial agglomeration surrounding the city of Padova are evident in the pronounced lobe of lichen desert running in a N-S

Station group n°	1	2	3	4
<i>Parmelia sulcata</i>	2	2	1	
<i>Evernia prunastri</i>	2	2	1	
<i>Hypogymnia physodes</i>	1	1	1	
<i>Parmelia exasperatula</i>	4	2	2	2
<i>Pseudevernia furfuracea</i>	1			
<i>Lecanora carpineae</i>	1	1		1
<i>Xanthoria fallax</i>		1		
<i>Usnea subfloridana</i>	1	1		
<i>Parmelia subrudecta</i>	1	1	1	
<i>Physconia distorta</i>		1		
<i>Lecanora allophana</i>	1			
<i>Lecanora conizaeoides</i>	1			
<i>Lecanora gr. hagenii</i>	1	1		1
<i>Arthonia radiata</i>	1	1		2
<i>Candelariella xanthostigma</i>	1	1	1	2
<i>Physcia biziana</i>		1		1
<i>Physcia vitii</i>	1	1	1	1
<i>Physconia grisea</i>	1	1		1
<i>Physcia adscendens</i>	1	1	1	3
<i>Hyperphyscia adglutinata</i>	1	1	1	2
<i>Lecanora chlorotera</i>	2	1	2	3
<i>Candelariella reflexa</i>	1	1	1	4
<i>Xanthoria parietina</i>	2	2	2	4
<i>Candelaria concolor</i>	2	1	2	5
<i>Lecidella elaeochroma</i>	4	2	2	3
<i>Physcia orbicularis</i>	3	2	5	5
<i>Physcia tenella</i>	5	5	5	5

Tab 5: Frequency vectors of the four station groups obtained by numerical classification of the matrix of species and stations relative to the province of Padova. Numbers refer to 5 frequency classes, with intervals of 20 % each.

direction, towards the province of Rovigo, which starts immediately south of Padova. A chain of hills (Colli Euganei) marks the western limit of this desert area; these hills act as a barrier to the dispersion of pollutants towards the southwest by the main winds (see fig.2); the barrier effect is evident in the southernmost portion of the province, where the absence of hills allows a much wider extension of the desert area towards the west. Considering the main wind directions shown in fig.2, it is probable that the pollutants emitted in the province of Venezia, and particularly in the large industrial area of Marghera, also contribute to the pollution stream flowing towards the province of Rovigo.






Fig. 14: I.A.P. map of the province of Padova. The province has been subdivided into several zones, characterized by different I.A.P. values.

Province of Vicenza

The province of Vicenza has a surface area of 2.722 km², and a population of 741.200 inhabitants, with a density of 238 inhabitants/km². The central and northern parts of the province (47.4% of the total surface), up to a line connecting Bassano and Schio, are below 100 m; its northern part (38.9 %) is occupied by a narrow belt of hills followed by a vast plateau with peaks reaching 1600 m. A small portion of the Lessinian Mountains occupies the central-western part of the province, west of Schio (13.7 % of the territory). The agricultural and industrial activities are very intense throughout the lowlands; the industries are generally scattered, with some main agglomerations around Vicenza. An important power station is located near Montorso, SW of Vicenza (see fig. 15).

This is the province with the highest number of recorded species (66). The most frequent are: *Physcia tenella* (85 % of the stations), *Physcia orbicularis* (82 %), *Candelaria concolor* (74 %), *Lecidella elaeochroma* (74 %), *Xanthoria parietina* (55 %) and *Candelariella reflexa* (52 %). This indicates a prevalence of *Xanthorion*-vegetation. The numerical classification is based on the 90 stations not reported in the preliminary survey (Nimis et al. 1991); it produced 7 main groups, which are reported in tab. 6. Group 1 includes only 4 stations, which differ markedly from all others in the high frequency of acidophytic species such as *Evernia prunastri*, *Hypogymnia physodes*, *Usnea* gr. *subfloridana*, *Pseudevernia furfuracea*, *Parmeliopsis ambigua*, *Scoliciosporum chlorococcum*, *Lecanora symmicta* etc., and in the low frequency of nitrophytic *Xanthorion* species: the relevés of this group, representing a *Parmelion*-vegetation, have all been taken on the hills north of Vicenza and on the Asiago Plateau, in a relatively undisturbed, pollution-free area. The group of stations 7 is also well characterized by the higher frequency of crustose lichens, such as *Lecanora carpinea*, *Caloplaca holocarpa*, *C. cerina*, *C. flavorubescens*, and *Lecanora hageni*, and represents a pioneer community leading towards *Xanthorion* vegetation. The other relevé groups have only slight floristic differences, and might be considered as different facies of *Xanthorion* vegetation.

As far as the distribution of the I.A.P. values is concerned (fig. 16), the province of Vicenza is characterized by marked differences. A true lichen desert is present only in the urban area of Vicenza, and close to an industrial area south of Thiene. Other areas with low I.A.P. values are located south of Bassano and near Montorso, in correspondence with two industrial zones. The highest I.A.P. values (more than 70) are reached on the Asiago Plateau and on the Lessinian Mountains. Also the areas lying east and south of Vicenza have relatively high I.A.P. values (30-50), which might be related to the prevailing northeastern winds. A much more detailed I.A.P. map of the Schio-Breganze area has been published by Nimis et al. (1991).

Station group n°	1	2	3	4	5	6	7
<i>Physconia grisea</i>	2	1	5	5	2	1	
<i>Hyperphyscia adglutinata</i>	2	3	5	4	1	1	3
<i>Parmelia exasperatula</i>	5	2	4		3	4	4
<i>Candelariella xanthostigma</i>	5	1	5	1	1	2	5
<i>Xanthoria parietina</i>	3	4	5	3	2	2	5
<i>Physcia adscendens</i>	4	2	4	2	1	2	5
<i>Physcia aipolia</i>	2	1	1		1	4	
<i>Lecanora chlorotera</i>	2	2	5	4	1	2	5
<i>Lecanora carpinea</i>	4	1	1			1	5
<i>Hypogymnia physodes</i>	4	1			1	2	2
<i>Evernia prunastri</i>	4			1	1	1	
<i>Usnea subfloridana</i>	5						1
<i>Parmelia sulcata</i>	5	1	2		1	3	5
<i>Chaenotheca furfuracea</i>	3	1			1	2	2
<i>Parmelia subaurifera</i>	2				1		
<i>Parmelia acetabulum</i>	2		1				
<i>Parmelia tiliacea</i>	3	1			1		
<i>Physconia distorta</i>	3	1		1	1	3	
<i>Caloplaca holocarpa</i>			1			4	
<i>Caloplaca cerina</i>					1	5	
<i>Lecanora gr. hagenii</i>		1	2		1	4	
<i>Ramalina fastigiata</i>	2						
<i>Lecidella flavosorediata</i>	2						
<i>Physcia stellaris</i>	3	1		1	1	4	
<i>Xanthoria fallax</i>	2	1	1		1	2	
<i>Physcia hirsuta</i>	2	1			1	1	
<i>Physcia biziana</i>		1	5	1	1	2	1
<i>Parmelia subrudecta</i>		1	1		1	1	
<i>Pseudevernia furfuracea</i>	4					1	1
<i>Parmeliopsis ambigua</i>	2						
<i>Bryoria furcellata</i>	2						
<i>Lecanora subfuscata</i>		1	1			1	
<i>Arthopyrenia punctiformis</i>	2	1		1	1	1	2
<i>Parmelia exasperata</i>			1			1	
<i>Normandina pulchella</i>						1	
<i>Graphis scripta</i>		1			1	1	
<i>Parmelia caperata</i>	2	1			1	1	
<i>Caloplaca flavorubescens</i>		1				2	
<i>Buellia punctata</i>						1	
<i>Physconia enteroxantha</i>						1	
<i>Physconia deterosa</i>						1	
<i>Scoliciosporum chlorococcum</i>	2						
<i>Lecanora symmicta</i>	2						
<i>Ramalina farinacea</i>					1		
<i>Pertusaria amara</i>						1	
<i>Parmelia saxatilis</i>					1		
<i>Parmelia quercina</i>				1			
<i>Opegrapha atra</i>		1					
<i>Heterodermia speciosa</i>						1	
<i>Cliostomum corrugatum</i>					1		
<i>Arthrosporum accline</i>		1		1			
<i>Catillaria nigroclavata</i>						1	
<i>Candelariella lutella</i>	2					3	
<i>Anaptychia ciliaris</i>						1	
<i>Parmelia subargentifera</i>	2	1				1	1
<i>Nectriaceae sp.</i>			1		1	1	1
<i>Parmelia glabra</i>						1	1
<i>Physcia vitii</i>	2	2	1		1	1	
<i>Arthopyrenia antecellens</i>		1	2	1	1	2	2
<i>Arthonia radiata</i>	3	2	3	1		1	2
<i>Rinodina exigua</i>		1	1		1	2	2
<i>Candelariella reflexa</i>	2	2	5	2	3	4	5
<i>Lecidella elaeochroma</i>	3	4	5	2	4	5	5
<i>Physcia orbicularis</i>	3	5	5		5	4	5
<i>Candelaria concolor</i>	3	5	5	4	2	4	5
<i>Physcia tenella</i>	4	5	5	4	5	5	4

Tab. 6: Table of species and station groups obtained by classification of the data relative to the province of Vicenza. Numbers refer to 5 frequency classes of 20% each.




Fig. 15: map of the province of Vicenza, with the location of the sampling stations.




Fig. 16: I.A.P. map of the province of Vicenza. The province has been subdivided into several zones, characterized by different I.A.P. values.

Province of Verona

The province of Verona has a surface area of 3.096 km², and a population of 741.244 inhabitants, with a density of 272 inhabitants/km². Most of the provincial territory (79.7 %) is occupied by lowlands, 16.1% by hills, and only 4.2% by mountains, located in the northern part of the province.

This is a species-rich province, with 56 species recorded. The most frequent ones are *Physcia tenella* (87 % of the stations), *Physcia orbicularis* (74 %), *Lecidella elaeochroma* (62 %), *Candelariella reflexa* (59 %), *Xanthoria parietina* (57 %), *Parmelia exasperatula* (51 %). With the exception of the latter, all of them are linked to *Xanthorion* vegetation. The numerical classification of the 118 stations produced 5 main groups, that are shown in tab. 7. Groups 1 and 2 have the highest species richness, and represent very well developed *Xanthorion* stands; the relevés have been taken in relatively undisturbed, pollution-free areas on the hills north of Verona; the group of stations 1 differs in the higher frequency of some *Parmelion*-species, such as *Parmelia quercina*, *P. tiliacea*, *P. sulcata*, *P. pastillifera*, *Pseudevernia furfuracea*, *Usnea subfloridana*, and represents a transition towards *Parmelion*-communities. The remaining three station groups are more species-poor, and show only slight compositional differences: they can be interpreted as three different facies of impoverished *Xanthorion* stands in more polluted areas; in particular, group 4 is characterized by the higher frequency of some acidophytic species, such as *Parmelia sulcata*, *Usnea subfloridana*, *Hypogymnia physodes*, *Lecanora conizaeoides*, *Parmelia exasperatula*, which indicates a secondary acidification of the bark due to air pollution, whereas group 5 is very species-poor, and the only constant species is *Physcia tenella*.

The I.A.P. map (fig. 18) shows wide differences within the provincial territory. The areas with the highest I.A.P. values (higher than 40 with single values reaching 70) are on the Lessinian Mountains, in the northern part of the province. An extended lichen desert occurs south of Verona; this city is located at the centre of a belt running in a E-W direction, characterized by I.A.P. values lower than 20; this belt includes the main highways connecting Verona with Vicenza eastwards, Mantova southwards and Brescia eastwards: along these highways there is a diffuse settlement of industrial activities. The dispersal of pollution east- and westwards from Verona agrees very well with the directions of the prevailing winds, as shown in fig. 2. The northern winds from the Adige Valley are responsible of the pronounced pollution lobe south of Verona, and the lower part of the Adige valley is also interested by pollution phenomena. Another belt with I.A.P. values mostly ranging between 20 and 30, separates this area from the southern part of the province where air pollution tends to worsen, with some areas of lichen desert occurring in its southernmost portion. In this case the pollution is allochthonous, deriving from sources located mainly in the province of Padova.

Station group n°	1	2	3	4	5
<i>Lecidella elaeochroma</i>	5	5	3	4	2
<i>Candelaria concolor</i>	5	5	4	2	1
<i>Physcia aipolia</i>	5	4	1	1	1
<i>Candelariella vitellina</i>	4	3	2	2	1
<i>Physcia stellaris</i>	4	2			
<i>Candelariella xanthostigma</i>	3	1	1	1	
<i>Rinodina exigua</i>	1	1	1	1	
<i>Physcia biziana</i>	2	2	2	2	1
<i>Physcia vitii</i>	1	1	1	2	
<i>Parmelia exasperata</i>	2	1			
<i>Parmelia quercina</i>	2	1			
<i>Parmelia tiliacea</i>	4	1	1	1	
<i>Parmelia subargentifera</i>	3	2			
<i>Parmelia acetabulum</i>	1	1		1	
<i>Parmelia sulcata</i>	4	1	1	2	1
<i>Parmelia glabra</i>	2				
<i>Arthopyrenia antecellens</i>	2	1			
<i>Caloplaca cerina</i>	3	2	1		
<i>Chaenotheca furfuracea</i>	1				
<i>Parmelia pastillifera</i>	2	1			
<i>Pseudevernia furfuracea</i>	2				
<i>Catillaria nigroclavata</i>	1				
<i>Nectriacea sp.</i>				1	
<i>Parmelia caperata</i>	1				
<i>Collema nigrescens</i>		1			
<i>Usnea subfloridana</i>	2		1		
<i>Scoliosporum chlorococcum</i>		1			
<i>Hypogymnia bitteriana</i>	1				
<i>Caloplaca ferruginea</i>	2	1			
<i>Parmelia glabratula</i>	1				
<i>Evernia prunastri</i>				1	1
<i>Lecanora subfuscata</i>	3	2	1	1	
<i>Hypogymnia physodes</i>	2	1	1	2	1
<i>Lecanora conizaeoides</i>				1	
<i>Parmelia subrudecta</i>	2	1	1	3	1
<i>Caloplaca flavorubescens</i>	1	1	1		
<i>Caloplaca holocarpa</i>			1	1	
<i>Arthonia radiata</i>	2	2	1		
<i>Physconia enteroxantha</i>	1				
<i>Xanthoria fallax</i>	3	2	1		1
<i>Physconia perisidiosa</i>	1				
<i>Physconia distorta</i>	3	2	1	1	1
<i>Lecanora allophana</i>	3	3	1	1	1
<i>Lecanora gr. hagenii</i>	3	3	2	2	1
<i>Parmelia exasperatula</i>	5	3	2	4	2
<i>Lecanora chlorotera</i>	5	4	2	2	1
<i>Lecanora carpinea</i>	5	4		1	1
<i>Physcia adscendens</i>	5	5	3	3	2
<i>Xanthoria parietina</i>	3	5	4	3	2
<i>Candelariella reflexa</i>	5	4	4	3	1
<i>Hyperphyscia adglutinata</i>	3	4	2	1	1
<i>Physconia grisea</i>	2	1	2	3	1
<i>Physcia tenella</i>	5	3	5	5	5
<i>Physcia orbicularis</i>	3	5	5	4	2

Tab 7: Table of species and station groups obtained by classification of the data relative to the Province of Verona.




Fig. 18: I.A.P. map of the province of Verona. The province has been subdivided into several zones, characterized by different I.A.P. values.

Province of Rovigo

The province of Rovigo occupies the southwestern portion of the region; it has a surface area of 1789 km², and a population of 250.000 inhabitants, with a density of 139 inhabitants/km².

All of the province is occupied by lowlands, and the provincial territory is mainly exploited by intensive agriculture. Natural or seminatural vegetation are present only in a few coastal sites and along the main watercourses.

This is the province where the lowest number of species has been recorded (23 species); the most frequent are *Physcia tenella* and *Lecidella elaeochroma*, that occur in more than 50 % of the stations, followed by *Parmelia exasperatula* (30%), *Physcia orbicularis* (25 %) and *Physcia adscendens* (25 %). Strongly nitrophytic and neutro-basiphytic species, such as *Candelaria concolor*, *Hyperphyscia adglutinata*, *Candelariella reflexa* and *Physconia grisea* are much less frequent in this province than in other lowland areas of the region with intense agricultural activity.

The classification of the 52 stations produced 5 main groups, reported in tab. 8: they all represent different, not very well characterized facies of *Xanthorion* vegetation, which indicates that the eutrophication of the bark due to agricultural activity occurs also in this province. However, the scattered occurrence of rather anitrophytic and acidophytic species, such as *Parmelia sulcata*, *P. exasperatula*, *Hypogymnia physodes*, *Usnea subfloridana* and *Lecanora conizaeoides*, the latter being more frequent within larger settlements, is an indication of the fact that the high pollution values reached in this province tend to produce a secondary acidification of the bark, counterbalancing the effects of eutrophication.. Of all the groups of stations, group 1 is characterized by a very low species frequency, and represents impoverished stands in very polluted areas.

The province of Rovigo has the worst air quality values within the region. Almost all of its territory has I.A.P. values lower than 20 (see fig. 20). The most polluted area (I.A.P. lower than 3, mostly lichen desert) is a 4-20 km wide belt, which crosses the surroundings of Rovigo in a SW-NE direction. Smaller lichen desert areas are located east of Sermide, in Emilia, and of Villanova Marchesana. A small area with I.A.P. higher than 20 is present in the northern coastal area, near Rosolina.

Near the coasts there is also one of the largest power stations of Italy. Considering the main directions of the winds, as shown by fig. 2 and by the general pollution map (fig. 21) it is probable that the effects of this power station are not particularly strong within the provincial territory. The main winds should transport the pollution in a southwestern direction, towards the region Emilia-Romagna (province of Ferrara), that borders the province of Rovigo to the east. A recent study on lichens as bioindicators in the province of Ferrara

Station group n°	1	2	3	4	5
<i>Parmelia tiliacea</i>			1		
<i>Candelariella xanthostigma</i>		1			1
<i>Candelaria concolor</i>					1
<i>Parmelia sulcata</i>	1	1	1		
<i>Arthonia radiata</i>	1				
<i>Usnea subfloridana</i>	1		1	1	
<i>Arthopyrenia antecellens</i>	1	1	1		
<i>Lecanora carpinea</i>	1			1	
<i>Lecanora gr. hagenii</i>		1		1	
<i>Parmelia exasperatula</i>	1	2	2	2	2
<i>Evernia prunastri</i>	1		1	1	2
<i>Hyperphyscia adglutinata</i>					1
<i>Candelariella reflexa</i>			1	1	
<i>Parmelia subrudecta</i>	1		1	3	2
<i>Hypogymnia physodes</i>	1		2	1	1
<i>Physconia grisea</i>			1		
<i>Xanthoria parietina</i>		2	1	1	3
<i>Lecanora chlorotera</i>		2		2	3
<i>Physcia orbicularis</i>	1	1	1	2	4
<i>Physcia adscendens</i>	2	2	2	1	2
<i>Lecanora conizaeoides</i>	1		2	1	1
<i>Lecidella elaeochroma</i>	5	2	2	5	4
<i>Physcia tenella</i>	1	5	3	5	5

Tab. 8: Frequency vectors of the five station groups obtained by numerical classification of the matrix of species and stations relative to the province of Rovigo. Numbers refer to 5 frequency classes, with intervals of 20 % each.

(Piccoli et al. 1989) shows that in this province there are high pollution levels, the lichen flora being represented by only a few species.

Since the main economic activity in this province is intensive agriculture, it is probable that most of the air pollution is allochthonous, deriving from the industrial areas of the provinces of Padova, Venezia and perhaps also Ferrara. This hypothesis is confirmed by the general pollution pattern at a regional scale (fig. 21) and by the main wind directions in the low plains (fig. 2). In the province of Rovigo the effects of phytotoxic pollutants may be further enhanced by the frequent persistence of fog during winter.




Fig. 19: map of the province of Rovigo, with the location of the sampling stations.




Fig. 20: I.A.P. map of the province of Belluno. The province has been subdivided into several zones, characterized by different I.A.P. values.




Fig. 21: I.A.P. map of the entire Region of Veneto. The regional territory has been subdivided into 7 zones, characterized by different I.A.P. values.

The I.A.P. map of the Region of Veneto

The I.A.P. map of the entire region is reported in fig. 21. The survey area has been subdivided into 7 zones, each with different pollution levels. The estimation of pollution levels by SO₂ is based on the correlations between direct pollution data and values of the I.A.P. index obtained in the study carried out at La Spezia (Nimis et al. 1990). In the following, each zone is briefly discussed.

Zone A (red colour) - air pollution: *high*. It includes the area with I.A.P. values ranging between 0 (lichen desert) and 5. From the results obtained at La Spezia these values should correspond to yearly mean values of the 98th percentiles of SO₂ higher than 84 µg/m³, yearly means of the 50th percentiles higher than 17 µg/m³, and winter means of the 50th percentiles higher than 27.7 µg/m³. In parts of this zone these values are probably higher, but cannot be revealed by the I.A.P. for the obvious reason that this zone includes also the lichen desert. This zone has a surface of 1.383 km², corresponding to 6.28 % of the entire region. The areas with the worst air quality include the surroundings of the industrial agglomeration of Marghera, a large part of the province of Padova, the greatest part of the province of Rovigo, and the surroundings of the city of Verona.

Zone B (bright orange) - air pollution: *moderate*. It includes the areas with I.A.P. values ranging from 6 to 10, corresponding to yearly mean values of the 98th percentiles of SO₂ ranging between 76 and 84 µg/m³, yearly means of the 50th percentiles ranging from 15.6 to 16.8 µg/m³, and winter means ranging from 25.2 to 27.6 µg/m³. This zone has a surface of 1624 km², corresponding to 7.45 % of the entire region. It tends to occupy a narrow belt all around the areas classified as zone A. It includes the centre of Treviso and its industrial area, a small industrial area in the eastern part of the province of Venice (near Portogruaro) and the centre of Vicenza.

Zone C (light orange) - air pollution: *rather low*. It includes areas with I.A.P. values ranging from 11 to 20, corresponding to yearly mean values of the 98th percentiles of SO₂ ranging from 61 and 75 µg/m³, yearly means of the 50th percentiles ranging from 13.2 and 15.5 µg/m³, and winter means ranging from 22.7 to 25.1 µg/m³. This zone has a surface of 3877 km², corresponding to 17.6 % of the entire region. It is particularly extended in the provinces of Venezia, Padova, Rovigo and Verona, it includes narrower areas in the provinces of Treviso and Vicenza, and the centre of the town of Belluno.

Zone D (yellow) - air pollution: *low*. It includes areas with I.A.P. values ranging from 21 to 30, corresponding to yearly mean values of the 98th percentiles of SO₂ ranging from 46 to 60 µg/m³, yearly means of the 50th percentiles ranging from 13.1 to 10.7 µg/m³, and winter means ranging from 17.7 to 22.6 µg/m³. This zone has a surface of 5150 km², corresponding to 23.4 % of the entire region. It extends over wide areas of the Venetian Plains, except their

eastern parts, and surrounds the areas included into the belts A-C, occupying also the southern part of the province of Belluno.

Zone E (light green) - air pollution: *very low*. It includes the areas with I.A.P. values ranging from 31 and 50, corresponding to yearly mean values of the 98th percentiles of SO₂ ranging from 17 to 45 µg/m³, yearly means of the 50th percentiles ranging from 5.8 and 10.6 µg/m³, and winter means ranging from 12.7 to 17.6 µg/m³. This zone has a surface area of 5917 km², corresponding to 28.9 % of the entire region. It occupies the largest part of the eastern section of the Venetian Plain, the outer part of the dealpine hills, and restricted portions of the provinces of Verona, Vicenza and Padova. It extends over a large portion of the province of Belluno.

Zone F (dark green) - air pollution: *negligible*. It includes the areas with I.A.P. values ranging between 51 and 70, corresponding to yearly mean values of the 98th percentiles of SO₂ ranging from 5 and 16 µg/m³, yearly means of the 50th percentiles ranging from 0.94 to 5.7 µg/m³, and winter means ranging from 2.71 to 12.6 µg/m³. This zone has a surface area of 3661 km², corresponding to 16.6 % of the entire region. It mainly extends over mountain areas, such as the Lessinian Mountains, the Asiago Plateau and the Dolomites in the province of Belluno.

Zone G (blue) - air pollution: *practically absent*. It includes areas with I.A.P. values higher than 70, corresponding to yearly mean values of the 98th percentiles of SO₂ lower than 5 µg/m³, yearly means of the 50th percentiles lower than 0.94 µg/m³, and winter means lower than 2.71 µg/m³. This zone occupies a narrow surface of 396 km², corresponding to 1.8 % of the region. It covers exclusively mountain areas, such as parts of the Lessinian Mountains, of the Asiago Plateau, and of the Dolomites.

In summary, the air pollution by SO₂ in the territory of the Veneto Region is as follows:

- in 6.3 % of the region there is a risk of high pollution levels (lichen desert areas).
- 25 % of the territory has a moderate air pollution (zones A-C).
- 23.38 % of the territory has a low air pollution (zone D).
- 44.93 % of the territory has a very low air pollution (zones E-G).

The southwestern part of the region has the highest pollution levels; it includes vast parts of the provinces of Rovigo, Padova and Venezia. The northern and eastern parts of the region have a better situation. By comparing the map of fig 21 with the direction of the prevailing winds (fig.2) it is evident that the pollution patterns shown by the map are well in accordance with the location of the main pollution sources (Mestre-Marghera, Padova, Verona etc.) and with the directions of the main winds. In particular, in the southern plains, the main transport is in a NE-SW direction, from Mestre towards the province of Rovigo.




Fig. 22: distribution map of *Candalaria concolor* in the Region of Veneto, obtained by automatic processing of its frequency data in the 662 stations.




Fig. 23: distribution map of *Candeleriella xanthostigma* in the Region of Veneto, obtained by automatic processing of its frequency data in the 662 stations.




Fig. 24: distribution map of *Hypogymnia physodes* in the Region of Veneto, obtained by automatic processing of its frequency data in the 662 stations.




Fig. 25: distribution map of *Parmelia exasperatula* in the Region of Veneto, obtained by automatic processing of its frequency data in the 662 stations.




Fig. 26: distribution map of *Parmelia sulcata* in the Region of Veneto, obtained by automatic processing of its frequency data in the 662 stations.




Fig. 27: distribution map of *Physcia adscendens* in the Region of Veneto, obtained by automatic processing of its frequency data in the 662 stations.




Fig. 28: distribution map of *Physcia orbicularis* in the Region of Veneto, obtained by automatic processing of its frequency data in the 662 stations.




Fig. 29: distribution map of *Xanthoria parietina* in the Region of Veneto, obtained by automatic processing of its frequency data in the 662 stations.

Distribution patterns of selected species

The matrix of the stations and species constitutes an interesting database for phytogeographic studies. For each species, data on its average frequency within each station are available. These can be processed by the same programs utilized to obtain the I.A.P. maps. The result is a series of distribution maps that, contrary to the currently used dot-maps, show not only the presence-absence of a given species within an area, but also its frequency distribution. It should be stressed that our data set does not refer to the complete distribution of the species within the survey area, since the substrate is limited to *Tilia*-boles. However, the restriction to one type of substrate only permits an easier interpretation of the maps, by excluding other edaphic factors that might influence the distribution of a given species. Distribution maps have been drawn for all species occurring in our data set. For reasons of space, in the following we report only the maps of eight selected species. They are briefly commented on in the following.

Candelaria concolor - This species, which is very common throughout Italy below the montane vegetation belt, is a typical element of *Xanthorion* vegetation. In north eastern Italy it has the optimum within the *Physcietum elaeinae* (Nimis & De Faveri 1981), a nitrophytic and heliophytic lichen community on eutrophic or eutrophiated bark, and it is often associated with *Hyperphyscia adglutinata* in areas with intense agricultural activities. According to Hawksworth & Rose (1970), when on eutrophic bark *Candelaria concolor* tolerates concentrations of SO_2 up to $40 \mu\text{g}/\text{m}^3$. Also Laundon (1973), De Wit (1976) and Deruelle (1983) cite this species as rather sensitive to air pollution in the British Isles, in Holland, and in northwestern France, respectively. Barkman (1958) attributes to this species a rather high poleophoby for the town of Oslo. In all of these cases the species is near to its northern distributional limits. A study by Sergio & Bento-Pereira (1981) demonstrates, on the contrary, that in Lisboa (Portugal) *Candelaria concolor* is one of the most tolerant lichens towards air pollution. A good resistance to air pollution has also been reported by Nimis (1985, 1986, 1990b) for the towns of Udine, Trieste and Rome.

The distribution map of *Candelaria concolor* in the Veneto Region (fig. 22) shows that this species has its optimum in the high plains, on the hills and on the lower mountains of the pre-Alps. It tends to decrease both in the relatively unpolluted mountain areas of the province of Belluno, and in the highly polluted parts of the southwestern plains. On a larger scale, this is the same behavior reported by Nimis et al. (1991) for the northern part of the province of Vicenza, and reflects both the moderately high tolerance of this species to air pollution, and its preference for eutrophiated substrata.

Candelariella xanthostigma - In Italy this species is widespread from the montane zone to the lowlands; in the South, however, it is mostly restricted to the mountains. There are only a few data on its tolerance to air pollution, particularly because it has been frequently confused with other congeneric species. Skye (1968) considers it as moderately sensitive to air pollution in Stockholm, and De Wit (1976) lists it under the rather pollution-intolerant species. Data from Italy (Nimis 1985, 1990b, Nimis et al. 1991) indicate that in this country *Candelariella xanthostigma* is rather poleophobic, and tends to be absent or rare within urban settlements and/or in heavily polluted areas.

The distribution map of *Candelariella xanthostigma* in the survey area (fig. 23) shows a distribution pattern similar to that of *Xanthoria parietina* and *Physcia adscendens*, with the difference that *Candelariella xanthostigma* seems to be more restricted to the relatively unpolluted mountain areas. It might also be that elevation plays a role in determining its distribution pattern within the region.

Hypogymnia physodes - This species is common throughout Italy from the lowlands (where it is less frequent) to the subalpine vegetation belt. It has a very wide ecological amplitude, being most frequent in communities growing on acid to subacid bark. According to Jürging (1975) its optimal pH is around 4.5. This species is one of the most frequently cited as an example of a pollution-tolerant lichen. The upper tolerance limits to SO₂ concentrations reported in the literature are 70-100 µg/m³ (Hawksworth & Rose 1970, British Isles), 75 µg/m³ (Türk & Christ 1986, Vienna), and 125-150 µg/m³ (Crespo et al. 1981, winter means, Spain). Data from Italy (Nimis 1985, 1986) show that this species can penetrate up to the limits of the lichen desert only in areas with a relatively humid climate, while in drier areas it seems to be rather poleophobic, probably because of the "drought effect", i.e. the drier microclimate of large urban areas.

The map of *Hypogymnia physodes* in the survey area (fig. 24) shows a disjunct distribution: the optimum is in the relatively unpolluted mountain areas with low agricultural activity; this species, which is mostly absent from the northern plain, reappears, with much lower frequency, around the most polluted areas of the low plains. This distribution seems to be conditioned essentially by the availability of acid substrates, which are present in the mountains (no or scarce eutrophication) and around the lichen desert areas (secondary acidification by pollution).

Parmelia exasperatula - This lichen, which is common throughout Italy, from the lowlands to the subalpine vegetation belt, has a wide ecological amplitude, being a common element of heliophytic lichen communities both on moderately eutrophiated and acidic bark. According to Jürging (1975) its pH optimum is around 5.2, and Wirth (1980) claims that it can stand a rather strong

eutrophication of the bark. According to Hawksworth & Rose (1970) *Parmelia exasperatula* does not tolerate mean concentrations of SO₂ higher than 50 µg/m³. De Wit (1976) lists this species as rather sensitive to air pollution in Holland and Skye (1968) claims that it is very sensitive in the surroundings of Stockholm. Data from Italy (Nimis 1985, 1986, Nimis et al. 1991) show that, at least in the north of the country, this species can penetrate up to the margins of the lichen desert.

The distribution map of *Parmelia exasperatula* in the Veneto Region (fig. 25) shows that its optimum is in the relatively unpolluted areas of the mountains. It tends to decrease both in the intensely cultivated high plains, and in the very polluted areas of the low plains. Also in this case, the behavior seen on a large scale very much resembles that reported by Nimis et al. (1991) for the northern part of the province of Vicenza. This distribution pattern is very similar to that of *Candelariella xanthostigma*.

Parmelia sulcata - This is one of the most widespread and common lichens world wide. In Italy it occurs from the lowlands to the subalpine belt, and is a frequent member of several lichen communities, mostly on subacid to acid bark. Its pH optimum, according to Jürging (1968) is of 5.2. In northern Italy it is rather photophytic at high elevations, rather skiophytic in the lowlands, where it is mostly confined at the northern side of isolated trunks. This species appears to be able to tolerate rather high pollution levels; the following upper limits are available in the literature: 80 µg/m³ (Seaward & Hitch, 1982, British Isles), from 60 to 125 µg/m³ (Crespo et al., winter means, Spain), 70 µg/m³ (Deruelle 1977, northern France). Data from northern Italy (Nimis 1985, 1986) indicate that *Parmelia sulcata* is able to thrive up to the limits of the lichen desert in areas with rather humid climate conditions.

The distribution map of *Parmelia sulcata* in the Veneto Region (fig. 26) shows a rather peculiar pattern: this species is most frequent in the relatively unpolluted mountain areas of the provinces of Belluno and Vicenza; it is absent both from the high plains and from most of the heavily polluted areas in the province of Rovigo, and it appears again, although with low frequency values, in the low plains, chiefly in the provinces of Treviso, Venice and Padova. A similar behavior has been reported by Nimis et al. (1991) for the northern part of the province of Vicenza. These distribution patterns might be explained as follows: *Parmelia sulcata* being an acidophytic and relatively anitrophytic lichen, has its optimum in the parts of the region where agricultural activity is lowest. It appears again in the plains where a moderately high pollution produces a secondary acidification of the bark.

Physcia adscendens - This species is widespread throughout Italy, from the lowlands to the subalpine, and sometimes even to the alpine vegetation belts; it

grows within *Xanthorion*-communities, and it appears to be somewhat less nitrophytic than *Physcia orbicularis*. Its tolerance to SO_2 is rather high: the upper tolerance limit reported in the literature is at $70 \mu\text{g}/\text{m}^3$ (Hawksworth & Rose 1970, Deruelle 1977). The average pH, according to Jürging (1975), is around 5.6. Data from Northern Italy (Nimis 1985, 1986, Nimis et al. 1991) indicate that *Physcia adscendens* is able to thrive up to the limits of the lichen desert, when it grows on eutrophic bark.

The distribution pattern of *Physcia adscendens* in the Veneto Region (fig. 27) is well correlated with the I.A.P. map (fig. 21). This species is most frequent in the relatively unpolluted province of Belluno, and progressively decreases in frequency towards the plains, being absent in the most polluted southwestern part of the region. This distribution pattern is very similar to that of *Xanthoria parietina* (see fig. 29), and will be discussed with the comments on this species.

Physcia orbicularis - This lichen is very common throughout Italy, from the lowlands to the montane vegetation belt. It grows in communities of the *Xanthorion parietinae*, and it is often associated with other nitrophytic species such as *Candelaria concolor* and *Hyperphyscia adglutinata* in intensely cultivated areas (Nimis & De Faveri 1981). In the British Isles *Physcia orbicularis* tolerates average concentrations of SO_2 up to $60 \mu\text{g}/\text{m}^3$, but it starts to become abundant only at values lower than $50 \mu\text{g}/\text{m}^3$ (Hawksworth & Rose 1970). Similar tolerance values are reported by Deruelle (1983) for the surroundings of Paris (France). Data from Italy (Nimis, 1985, 1986, 1990b, Nimis et al. 1991) indicate that this species can thrive up to the limits of the lichen desert, particularly on eutrophic or eutrophied bark.

In the Veneto Region *Physcia orbicularis* is one of the commonest epiphytic species, and one of those that are more able to stand high air pollution levels. Its distribution map (fig. 28) shows a pattern that very much resembles that of *Candelaria concolor*. *Physcia orbicularis* has its optimum in the high plains, the hills and the pre-Alps, becoming progressively less frequent in the most polluted areas around large settlements and in the provinces of Rovigo and Padova. Its distribution seems to be influenced positively by the secondary eutrophication of the barks due to the agricultural activity, and negatively by strong air pollution levels.

Xanthoria parietina - This species is very common throughout Italy, from the lowlands to the upper montane vegetation belt; it has its optimum within *Xanthorion*-vegetation, and it is somewhat less nitrophytic than e.g. *Physcia orbicularis* and *Candelaria concolor*. Its optimal pH, according to Jürging (1975) is of 5.6. The upper tolerance limits to average concentrations of SO_2 available in the literature are: $70 \mu\text{g}/\text{m}^3$ (Hawksworth & Rose 1970, British Isles), $80\text{-}90 \mu\text{g}/\text{m}^3$ (Johnsen & Söchting 1973, Copenhagen). Apparently, this species is

rather tolerant to pollution, although the reports on its poleophoby throughout Europe are rather contradictory (see Nimis et al. 1991). Data from Italy (Nimis 1985, 1986, 1990b, Nimis et al. 1990, 1991) indicate a moderate to rather high toxitolerance for *Xanthoria parietina*.

The distribution pattern of *Xanthoria parietina* in the survey area (fig. 29) very much resembles that of *Physcia adscendens* (fig. 27); the two species also have a similar ecology and are frequently associated. The distribution of both species is well in agreement with the I.A.P. map (fig 21): both have frequency maxima in the relatively unpolluted mountain areas, and decrease progressively towards the most polluted parts of the southwestern plains. Their distribution is probably influenced, negatively, both by air pollution and by the strong eutrophication of the bark due to agricultural activities.

The distribution maps shown in figs 22-29 can be classified into 4 main groups, that represent 4 main distribution patterns. Each group is briefly discussed below.

Distribution pattern 1). - *Candelariella xanthostigma*, *Parmelia exasperatula*.

These are slightly nitrophytic or anitrophytic, subacidophytic to subneutrophytic, rather pollution-intolerant species. Their distribution in the survey area is limited both by air pollution and by the secondary eutrophication of the bark. Their optimum is in the relatively unpolluted mountain areas with low agricultural activity.

Other species with a similar distribution pattern are: *Arthopyrenia antecellens*, *Lecanora chlarotera*, *Lecanora argentata*, *Parmelia caperata*, *Parmelia subaurifera*, *Parmelia tiliacea*, *Parmeliopsis ambigua*.

Distribution pattern 2). - *Physcia adscendens*, *Xanthoria parietina*.

These are rather nitrophytic, subneutrophytic, rather pollution-tolerant species. Their distribution pattern is intermediate between those of group 1) and of group 3), since they are less sensitive to air pollution than the species of group 1, and more sensible to strong eutrophication than the species of group 3.

Other species with a similar distribution are: *Caloplaca cerina*, *Parmelia acetabulum*, *Parmelia subrudecta*, *Physcia aipolia*, *Physconia distorta*, *Xanthoria fallax*.

Distribution pattern 3) - *Candelaria concolor*, *Physcia orbicularis*.

These are strongly nitrophytic, neutro-basiphytic, pollution-tolerant species. Their distribution is limited by pollution in the southwestern part of the region, by scarce eutrophication of the bark in the northern part. Their optimum is in the intensely cultivated, but less polluted northern plains, the hill belt and the parts of the pre-Alpine valleys with rather high agricultural activity.

Other species with a similar distribution pattern are *Candelariella reflexa*, *Hyperphyscia adglutinata* and *Physconia grisea*.

Distribution pattern 4) - *Parmelia sulcata*, *Hypogymnia physodes*.

These are acidophytic, scarcely or non- nitrophytic, rather or strongly pollution-tolerant species. The main limiting factor for these lichens is the availability of acid to subacid, non- or scarcely eutrophicated substrates. These are available in the mountain areas with scarce agricultural activity and in the areas surrounding the lichen desert, where secondary acidification by SO₂ is stronger. This explains their disjunct distribution patterns, with an optimum in the mountains, and a scarce presence also around the most polluted areas of the plains.

Other species with a similar distribution pattern are: *Chaenotheca furfuracea*, *Evernia prunastri*, *Lecanora conizaeoides*, *Pseudevernia furfuracea*, *Usnea* gr. *subfloridana*.

Noteworthy is the good correspondence between the ecology of all cited species (e.g. as evident in the ordinations of figs. 5 and 6) and their distribution patterns.

The distribution of lichens within the region is conditioned not only by air pollution by SO₂, but also by other factors, the most important of which seems to be the secondary eutrophication of the bark due to the agricultural activity.

CONCLUDING REMARKS

The results of this study allowed the production of the first large-scale air pollution map based on bioindicators in Italy. The method adopted in this study proved to have a significant correlation with pollution values; the pollution patterns shown by the maps are well in agreement with the location of the main emitting sources and with the directions of the prevailing winds. The method allowed also the detection of a widespread and pronounced pollution by inorganic and organic fertilizers, the eutrophication of the bark tending to counterbalance the acidifying effects of pollution by sulphur dioxide. The maps presented in this paper may be utilized to detect high risk areas within the region, and to optimize the positioning of the recording gauges. Considering that this study has been carried out within one year, and that its costs were lower than the cost of a single recording gauge, the joint use of bioindicators and of instrumental recording is suggested as one of the most appropriate approaches to air pollution monitoring.

Acknowledgements

The authors are grateful to Drs. E. Cosma and S. Olivieri (Region of Veneto), for the comments to the manuscript and for help during field work, and to Drs. M.G. Bellio, G. Bolognini, M. Castello, K. Kravos, M. Tretiach (Trieste), for assistance in data elaboration.

RIASSUNTO - Vengono presentati i risultati di uno studio sull'inquinamento da SO₂ nella regione del Veneto, effettuato utilizzando i licheni epifiti come bioindicatori. E' stato adottato il metodo basato sull' "Indice Lichenico Calibrato di Qualità dell' Aria" proposto da Liebenoerfer et al. (1988). Tale indice viene calcolato sommando le frequenze di tutte le specie di licheni epifiti all'interno di un reticolo di rilevamento composto da 10 unità, ed ha dimostrato una correlazione altamente significativa con la somma delle concentrazioni di diverse sostanze inquinanti. I risultati di due studi preliminari condotti dagli autori in territorio italiano, a La Spezia e nell'Alto Vicentino hanno permesso di confermare l'alta predittività di tale indice per l'inquinamento da anidride solforosa.

Nel complesso, sono stati effettuati 2425 rilevamenti in 662 stazioni, così distribuite tra le varie provincie:

Provincia di Belluno: 30 stazioni

Provincia di Padova: 80 stazioni

Provincia di Rovigo: 65 stazioni

Provincia di Treviso: 80 stazioni

Provincia di Venezia: 60 stazioni

Provincia di Verona: 120 stazioni

Provincia di Vicenza 227 stazioni.

L'alto numero di stazioni in provincia di Vicenza deriva dal fatto che nella sua parte settentrionale è stato condotto uno studio preliminare per saggiare le metodiche successivamente estese all'intero territorio regionale.

Tutti i rilievi di vegetazione epifita sono stati effettuati su Tiglio. I rilievi sono stati effettuati utilizzando una griglia di 30 x 50 cm, suddivisa in 10 rettangoli, sistemata ad 1.2 m sul tronco, con il centro nell'area con la massima copertura lichenica. La somma delle frequenze di tutte le specie presenti nei 10 rettangoli costituisce l' Indice di Purezza Atmosferica (I.A.P.) del rilievo, L' I.A.P. della stazione è la media dei valori di tutti i rilievi effettuati nella stazione stessa (in media 3.7).

Le matrici delle specie e delle stazioni sono state sottoposte ad elaborazione tramite programmi di analisi multivariata (classificazione ed ordinamento) al fine di individuare i principali aspetti vegetazionali in ciascuna provincia, ed i principali gradienti ecologici sottostanti alla variazione floristica nell'ambito dell'intera regione.

L' elaborazione cartografica dei dati è stata effettuata tramite il programma SURFER (Goden Software Inc., Golden, colorado, U.S.A.). I dati di input sono: a) dati di I.A.P. delle stazioni per le sette provincie e per l'intero territorio regionale (carte

dell'inquinamento); b) frequenza media di ciascuna specie (media delle frequenze dei rilievi di ciascuna stazione) per tutte le stazioni (carte di distribuzione di specie singole).

Vengono presentate le carte dell' inquinamento da SO₂ delle sette provincie, e dell'intera regione: esse sono in buon accordo sia con la disposizione delle principali fonti emittenti che con la direzione prevalente dei venti. Nel 93.7 % del territorio regionale le medie annue dei 98ⁱ percentili di SO₂ sono inferiori a 84 µg/m³, e quindi il rischio di soglie di inquinamento superiori ai limiti stabiliti dalla legge è nullo o molto basso. Il 25 % del territorio regionale presenta un moderato inquinamento atmosferico, il 23.4 % un inquinamento basso, il 44.9 % ha livelli di inquinamento da SO₂ trascurabili. Le aree maggiormente inquinate si trovano nella porzione sudoccidentale del Veneto, nelle provincie di Padova, Rovigo e Vicenza. Ulteriori aree ad inquinamento maggiore sono site presso altri centri urbani ed aree industriali. La provincia meno inquinata è quella di Belluno, quella più inquinata è quella di Rovigo, ove tuttavia l'inquinamento proviene in gran parte da aree site al di fuori del territorio provinciale, soprattutto dalle provincie di Padova e Venezia. Le carte possono venir utilizzate per individuare aree a rischio in cui posizionare le centraline per il rilevamento strumentale dell'inquinamento.

La vegetazione lichenica prevalente nell'ambito degli alberi campionati in questo studio è quella appartenente all'alleanza *Xanthorion parietinae*, che include comunità di licheni fotofili, nitrofilo, neutro-basofili. Ciò è dovuto in primo luogo all'eutrofizzazione secondaria dei substrati conseguente all'intensa attività agricola nella Pianura Veneta.

Vengono presentate le carte di distribuzione, anch'esse ottenute per via automatica, di 8 specie: *Candelaria concolor*, *Candelariella xanthostigma*, *Hypogymnia physodes*, *Parmelia exasperatula*, *Parmelia sulcata*, *Physcia adscendens*, *Physcia orbicularis*, *Xanthoria parietina*. Esse mostrano un' ottima concordanza tra patterns di distribuzione ed ecologia delle specie stesse. In particolare, si possono distinguere 4 patterns di distribuzione principali: 1) specie acidofitiche e non nitrofitiche, sensibili all'inquinamento, con optimum nelle aree montane a scarso sviluppo agricolo; la loro distribuzione è limitata sia dall'inquinamento atmosferico da anidride solforosa che da quello da fertilizzanti organici ed inorganici; 2) specie moderatamente nitrofitiche, subneutrofitiche, moderatamente tolleranti rispetto all'inquinamento da anidride solforosa; queste mostrano dei pattern di distribuzione intermedi tra quelli del gruppo precedente e del gruppo seguente; 3) specie fortemente nitrofitiche, neutro-basifitiche, resistenti all'inquinamento da anidride solforosa; il loro optimum è nell'alta Pianura Veneta e nella zona prealpina con maggiore sviluppo agricolo; la loro distribuzione è limitata dall'inquinamento da anidride solforosa nella parte sudoccidentale della regione, dalla scarsa eutrofizzazione dei substrati in quella montana; 4) specie relativamente anitrofitiche ed acidofitiche, che presentano un optimum di distribuzione nelle aree montane a scarso sfruttamento agricolo, una scarsissima frequenza nelle parti della pianura veneta moderatamente o poco inquinate, ed un lieve aumento della frequenza attorno alle aree più inquinate della pianura; tale gruppo di specie evidenzia l' acidificazione secondaria dei substrati nelle aree maggiormente inquinate dovuta all' inquinamento da anidride solforosa.

I fattori principali che condizionano la presenza dei licheni nella Regione del Veneto sono l'inquinamento atmosferico da gas fitotossici e quello da fertilizzanti organici ed inorganici (eutrofizzazione delle scorze) conseguente all'intensa attività agricola.

References

- Ammann K., Herzig R., Liebendoerfer L. & M. Urech, 1989. *Statistical correlation of deposition data on 8 different air pollutants to Swiss lichen IAP method in Biel and the Swiss plain*. Manuscr.(in press).
- Barkman J.J., 1958. *Phytosociology and ecology of cryptogamic epiphytes*. Van Gorcum and Co., Assen. 628 pp.
- Barkman J.J., 1969. *The influence of air pollution on bryophytes and lichens*. In: Air Pollution, Proceedings of the 1st European Congress on the Influence of Air Pollution on Plants and Animals. Wageningen, 1968:237-241. Centre for Agr. Publ. and Doc.Wageningen.
- Beschel R.E. 1958. *Flechtenvereine der Städte, Stadtflechten und ihr Wachstum*. Ber. Naturw.-med. Ver. Innsbruck, 52: 1-158.
- Caniglia G., Giulini P. & M. Spampiani, 1978. *Inquinamento atmosferico e licheni. Saggio di distribuzione nella Valle del Boite e a Cortina d'Ampezzo*. Atti IV Conv. Ecologia Alpi Orientali, Pian Cansiglio:279-293.
- Coppins B.J. 1973. *The "Drought Hypothesis"*. In Ferry et al., op.cit.: 124-142.
- Crespo A., Barreno E., Sancho L.G. & A.G. Bueno, 1981. *Establecimiento de una red de valoración de pureza atmosférica en la provincia de la Coruña (España) mediante bioindicadores liquenicos*. Lazaroa, 3: 289-311.
- Del Favero R., O. Andrich, G. Dal Mas, C. Lasen & L. Poldini, 1990. *La Vegetazione forestale del Veneto. Prodrumi di tipologia forestale*. Regione del Veneto. Venezia. 180 pp.
- Deruelle S., 1977. *Influence de la pollution atmosphérique sur la végétation lichénique des arbres isolés dans la région de Mantes (Yvelines)*. Rev. Bryol. Lichénol., 43 (2): 137-158.
- Deruelle S., 1978. *Etude comparée de la sensibilité de trois méthodes d'estimation de la pollution atmosphérique, en utilisant les lichens comme indicateurs biologiques dans la région de Mantes (Yvelines)*. Rev. Bryol. Lichénol., 44 (4): 429-441.
- Deruelle S., 1983. *Écologie des lichens du bassin Parisien. Impact de la pollution atmosphérique et relations avec les facteurs climatiques*. Univ. P. M. Curie. These de Doct. 2 voll.
- De Wit T., 1976. *Epiphytic lichens and air pollution in the Netherlands*. Bibl. Lichenol., Band 5, Cramer, Vaduz. 228 pp.
- Ferry B.W., Baddeley M.S. & D.L. Hawksworth (eds.), 1973. *Air Pollution and Lichens*. University of Toronto Press. Toronto. 390 pp.
- Hawksworth D.L. & L. Rose, 1970. *Qualitative scale for estimating sulphur dioxide air pollution in England and Wales using epiphytic lichens*. Nature, 227: 145-148.
- Herzig R., 1990. *Entwicklung des Integrierten biologischen Messsystems der Luftverschmutzung mit Flechten in der Schweiz, seine Anwendung und erste Vergleiche mit Bodenanalysen*. VDI Berichte 837: 937-956.

- Herzig R., Liebendoerfer L. & M. Urech, 1985. *Flechten als biologische Indikatoren der Luftverschmutzung in der Schweiz. Methodenentwicklung in der Region Biel-Seeland*. Lizentiatsarbeit Univ. Bern, 241 pp.
- Herzig R., Liebendoerfer L., M. Urech & K. Ammann, 1987. *Evaluation und Kalibrierung einer Flechtenindikationsmethode mit wichtigen Luftschaden*. In: Bioindikation - Wirkungsbezogene Erhebungsverfahren für den Immissionschutz. Freising VDI-Koll., 609: 619-639.
- Johnsen I. & U. Söchting, 1973. *Influence of air pollution on the epiphytic lichen vegetation and bark properties of deciduous trees in the Copenhagen area*. Oikos, 24: 344-351.
- Jürging P., 1975. *Epiphytische Flechten als Bioindikatoren der Luftverunreinigung dargestellt an Untersuchungen und Beobachtung in Bayern*. Bibl. Lichenol., 4, 164 pp.
- Lackovicova A., Lisická E. & I. Pisut, 1988. *Literature on lichen indication in socialist countries*. Biologia (Bratislava), 43, 1: 79-86.
- Lackovicova A., Lisická E. & I. Pisut, 1989. *Literature on lichen indication in socialist countries*. 2. Biologia (Bratislava), 44, 1: 83-89.
- Laundon J.R., 1973. *Urban Lichen Studies*. In: Ferry B.W., Baddeley M.S. & D.L. Hawksworth (eds.), op. cit.: 109-123.
- Liebendoerfer L., Herzig R., Urech M., & K. Ammann, 1988. *Evaluation und Kalibrierung der Schweizer-Indikationsmethode mit wichtigen Luftschadstoffen*. Staub - Reinhaltung der Luft, 48: 233-238.
- Nimis P.L., 1985. *Urban Lichen Studies in Italy. I: the town of Trieste*. Studia Geobot., 5: 49-74.
- Nimis P.L., 1986. *Urban Lichen Studies in Italy. II: the town of Udine*. Gortania, 7 : 147-172.
- Nimis P.L., 1990. *Air Quality indicator and indices: the use of plants as bioindicators for monitoring air pollution*. Proc. Int. Workshop on Indicator and Indices JRC Ispra, EUR 13060 EN: 93-126.
- Nimis P.L., 1990b. *Urban Lichen Studies in Italy. III: the city of Rome*. Annali di Botanica (in press).
- Nimis P.L., Castello M. & M. Perotti, 1990. *Lichens as biomonitors of sulphur dioxide pollution in La Spezia (Northern Italy)*. Lichenologist, 22 (3): 333-344.
- Nimis P.L., Bargagli R., Benedet A., Castello M., Gasparo D., Lazzarin G. & A., Olivieri S. & M. Tretiach, 1991. *Licheni come bioindicatori di inquinamento atmosferico nell'Alto Vicentino*. Atti Mus. St. Nat. Verona (in press).
- Nimis P.L. & R. De Faveri, 1981. *A numerical classification of Xanthorion communities in NE Italy*. Gortania, 2: 91-110.
- Nimis P.L. & J. Poelt, 1987. *The lichens and lichenicolous fungi of Sardinia. An annotated list*. Studia Geobot., 7, suppl. 1, 269 pp.
- Piccoli F., Kumer E. & L. Bonalberti, 1989. *Licheni ed inquinamento atmosferico a Ferrara*. Archivio Botanico Italiano, 65 (1-2): 73-80.
- Rydzak J., 1969. *Lichens as indicators of the ecological conditions of the habitat*. Annls. Univ. Mariae Curiae Sklodowska, C, 23: 131-164.

- Seaward M.R.D. & J.C.B. Hitch (eds.), 1982. *Atlas of the lichens of the British Isles*. Univ. of Bradford.
- Sergio C. & F. Bento-Pereira, 1981. *Liquenes e briofitos como bioindicadores da poluição atmosférica. I. Utilização de uma escala qualitativa para Lisboa*. Bol. Soc. Brot. Ser.2, 54: 291-303.
- Skye E., 1968. *Lichens and air pollution. A study of cryptogamic epiphytes and environment in the Stockholm region*. Acta Phytogeogr. Suec., 52: 1-123.
- Spampani M., 1982. *I licheni: indicatori fisiologici della qualità dell'aria*. Le Scienze, 167: 60-69.
- Türk R. & R. Christ, 1986. *Beitrag zur epiphytischen Flechtenflora im Stadtgebiet von Wien*. Verh. zool.bot. Ges. Oesterr., 124: 65-80.
- Urech M., Peter K., Liebendoerfer L. & R. Herzig. 1990. *Bioindikation mit Flechten im Kanton Luzern*. Mitt. Naturforsch. Ges. Luzern, 31: 219-232.
- Walter H. & H. Lieth, 1960. *Klimadiagramm-Weltatlas*. Jena.- Fischer Verlag.
- Wildi O. & L. Orloci, 1983. *Management and multivariate analysis of vegetation data*. 2nd ed. Swiss Fed. Inst. For. Res., Rep. 215. Birmensdorf. 387 pp.
- Wirth V., 1980. *Flechtenflora*. Ulmer. Stuttgart, 552 pp.

Authors' Addresses

Prof. Pier Luigi NIMIS
Dipartimento di Biologia
Università di Trieste
Via A. Valerio 32
I 34127 TRIESTE

Dr. Arianna LAZZARIN
Cooperativa Co.Ge.V.
Lungadige Sarmicheli 9
I 37100 VERONA

Dr. Giuliano LAZZARIN
Museo Civico di Storia Naturale di Verona
Sezione Botanica
C.so Cavour 11
I 37121 VERONA

Dr. Dario GASPARO
Cooperativa ECOTHEMA
Via Stuparich 15
I 34100 TRIESTE

Studio fitosociologico e cartografia della vegetazione della Sila Greca (Calabria)

Liliana BERNARDO, Giuliano CESCO, Michele CODOGNO,
Simona FASCETTI e Domenico PUNTILLO *

Keywords: mapping, phytosociology, Sila Greca (Calabria, S-Italy), vegetation.

Abstract. *PHYTOSOCIOLOGICAL STUDY AND MAP OF THE VEGETATION OF SILA GRECA (CALABRIA, S-ITALY).* This paper presents the results of a study on the vegetation of the Sila Greca. The enclosed map (1:50,000) is based on the physiognomy and structure of vegetation. The floristic composition of the formation-types represented in the vegetation map is described in the text.

Introduzione

Questo studio e la carta vegetazionale annessa sono stati realizzati nell'ambito di un Progetto Finalizzato del Consiglio Nazionale delle Ricerche, avente per scopo lo studio multidisciplinare dei sistemi agricoli e forestali, con particolare riguardo alle aree marginali. Una delle aree campione scelte per tale ricerca è stato il territorio della Comunità Montana della Sila Greca.

La Sila Greca è una zona ancora poco nota sotto il profilo botanico, per cui un documento di sintesi cartografica, che fornisce un quadro fitosociologico di tale area, può in parte colmare le lacune esistenti a tale riguardo.

Ambiente fisico

La Sila Greca è situata sul versante ionico del massiccio silano nel tratto esposto a NE e compreso fra i corsi del torrente Cino e del fiume Nicà (Fig. 1). Essa è delimitata a nord dal Mare Ionio ed a sud da una serie di rilievi (M. Altare, 1653 m; M. Paleparto, 1480 m), che la separano dalla Sila Grande. Questa regione è caratterizzata da un elevato grado di acclività: si raggiungono 1500 m di altitudine a solo 20 km dalla costa.

* Il presente lavoro è stato finanziato con fondi C.N.R., Progetto Finalizzato "Incremento della Produttività delle Risorse Agricole" (resp. Prof. G. Cesca) e con fondi M.P.I. 60 % (resp. Dr. M. Codogno).




Fig. 1 - Carta schematica della Sila Greca. I tipi climatici indicati seguono la classificazione di Thornthwaite (1948).
Schematic map of the Sila Greca. The climate-types follow the Thornthwaite's classification (1948).

Geologia e morfologia

Il substrato geologico della Sila Greca è costituito da un complicato alternarsi di litotipi diversi, ascrivibili a quattro complessi principali (Puglisi *et al.*, 1985). Il complesso più antico e più esteso è quello che forma la parte interna dell'area studiata ed è costituito da rocce ignee intrusive acide (graniti) e metamorfiche (gneiss, filladi, ecc.). Il secondo complesso, trasgressivo sul primo, è composto in prevalenza da rocce calcaree (serie di Longobucco), costituenti la copertura mesozoico-terziaria dello zoccolo calabrese. Il terzo complesso è costituito da rocce sedimentarie (arenarie, marne, argille, ecc.), la cui deposizione risale ad un periodo che va dall'Oligocene al Pliocene. Il quarto complesso infine è composto dai depositi alluvionali di origine quaternaria.

La morfologia del territorio è strettamente condizionata dall'associazione di unità litologiche con gradi di plasticità e resistenza all'erosione molto diversi. L'attuale situazione geomorfologica può essere schematizzata individuando tre zone all'interno delle quali i differenti litotipi mostrano forme di modellamento affini:

- Altopiano silano al di sopra di 800-1000 m di altitudine: l'alterazione superficiale delle rocce (in prevalenza graniti) dà luogo alla formazione di uno strato di sabbia a tessitura grossolana. Lo spessore di questo mantello di alterazione è notevole nelle zone pianeggianti e sommitali a morfologia dolce, mentre sui versanti l'asporto continuo di materiale degradato determina l'affioramento della roccia madre.
- Sinistra orografica del medio bacino del Trionto: la morfologia di questo settore è caratterizzata dalla presenza di versanti ripidi e di valli molto strette e profonde, modellate sulle rocce appartenenti alla serie di Longobucco.
- Destra orografica del medio e basso bacino del Trionto: questa zona è costituita da terrazzamenti scavati in rocce sedimentarie di vario tipo (argille, sabbie, marne) dall'azione erosiva del mare durante il sollevamento isostatico di questa regione. Il substrato geologico, dove la copertura vegetale è scarsa e la pendenza è elevata, è soggetto a fenomeni di erosione accelerata, con formazione di frane e di calanchi.

Tutta la regione studiata è percorsa da corsi d'acqua a carattere periodico ("fiumare"), che incidono valli brevi con versanti simmetrici, forte inclinazione e tratto inferiore molto ampio e sovralluvionato.

Clima

Il macroclima della Sila Greca risulta essere di tipo mediterraneo, caratterizzato dall'alternarsi di una stagione invernale mite e piovosa e di una stagione estiva secca. L'ampio campo di variazione dell'altitudine (ca. 1500 m) porta però a notevoli variazioni topoclimatiche nell'area in oggetto (Fig. 2): infatti con l'aumentare della quota aumentano le precipitazioni (ca. 90 mm ogni




Fig. 2 - Climadiagrammi di tre stazioni meteorologiche nell'area di studio, poste a diverse altitudini.
Climatic diagrams of three weather-stations in the study area at different altitudes.

100 m) e diminuiscono le temperature (0,7-0,8°C ogni 100 m), con conseguente diminuzione dell'evapotraspirazione potenziale.

I dati termici e pluviometrici di 21 stazioni meteorologiche (Ciancio, 1971) sono stati elaborati secondo il metodo di Thornthwaite (1948) con lo scopo di calcolare l'evapotraspirazione potenziale e di individuare i limiti territoriali dei diversi tipi climatici per l'area studiata. Tali limiti sono stati riportati in Fig. 1. Questa figura mette in evidenza come dal clima di tipo semiarido della zona costiera si passi rapidamente al clima di tipo perumido delle quote più elevate.

Vegetazione

Dati e metodi

I tipi vegetazionali impiegati nella cartografia sono stati definiti mediante l'analisi di 145 rilievi fitosociologici eseguiti secondo il metodo di Braun-Blanquet (1964). L'individuazione dei limiti dei tipi vegetazionali è stata realizzata mediante l'interpretazione di una serie di foto aeree eseguite da bassa quota, messe a disposizione dalla Comunità Montana della Sila Greca.

Lo schema della carta è articolato in una classificazione principale della vegetazione secondo i piani altitudinali. Per la complessità della zonazione della vegetazione in senso verticale, la nomenclatura dei piani altitudinali impiegata è stata scelta con lo scopo di fornire indicazioni il più possibile generali sulle loro caratteristiche altitudinali ed ambientali. Per ogni piano altitudinale è stata eseguita una classificazione basata su caratteri fisionomici e strutturali in gruppi di formazioni e, nell'ambito di questi, una suddivisione su base principalmente floristica. Per tale scopo, le tabelle fitosociologiche ottenute nell'ambito dei diversi gruppi di formazioni sono state elaborate mediante programmi di analisi multivariata (Wildi e Orlóci, 1983) e di analisi nodale (Češka e Roemer, 1971).

La scelta della scala 1:50.000 è stata motivata da due ragioni: l'estensione del territorio e la presenza di vaste superfici ricoperte da vegetazione fisionomicamente omogenea. In tal senso, questa scala risulta la più opportuna in quanto fornisce un documento di sufficiente dettaglio e, nello stesso tempo, sintetizza l'attuale situazione della copertura vegetale del territorio.

La nomenclatura utilizzata per le specie vegetali è quella della *Flora d'Italia* del Pignatti (1982a).

Note fitogeografiche

Codogno e Puntillo (1987), in uno studio sulla fitogeografia dei prati e pascoli della Sila Greca, hanno messo in evidenza che i geoelementi maggiormente rappresentati nella composizione floristica delle formazioni erbacee di bassa quota sono quelli mediterranei con baricentro orientale (E-

eurimediterraneo e NE-stenomediterraneo), mentre in quelle di quota più elevata l'elemento orientale (SE-europeo) ed occidentale (W-europeo) coesistono. Ciò avviene pure per le associazioni ad arbusti spinosi della Sila, dove *Astragalus calabrus* (di origine orientale) e *Genista anglica* (di origine occidentale) coesistono nella stessa fascia altitudinale (montana). La composizione floristica della vegetazione forestale della Sila Greca presenta dal punto di vista corologico un comportamento analogo a quello succitato. Ciò è in perfetto accordo con la divisione della penisola appenninica in zone biogeografiche su base fitosociologica proposta da Bonin (1978, cartina a pag. 290), nella quale il territorio della Sila Greca risulta essere a cavallo fra la zona ionica (con affinità balcanico-meridionale) e quella tirrenica (più povera in elementi balcanici e decisamente a carattere più oceanico).

Zonazione altitudinale

La variazione della durata del periodo arido estivo con l'altitudine, determinata dalle diverse condizioni topoclimatiche succitate, dà luogo ad una zonazione altitudinale della vegetazione nel territorio studiato (Codogno e Puntillo, 1988). Per l'interpretazione di questa zonazione è necessario non trascurare l'azione antropica, la quale sia nel passato che attualmente ha apportato notevoli modificazioni alla vegetazione potenziale nell'ambito delle diverse fasce altitudinali. Tale zonazione può venir schematizzata come segue:

- *Piano basale*: si estende dalla linea di costa fino a 600 m di altitudine. La vegetazione zonale di questo piano è costituita da formazioni a sclerofille sempreverdi termoxerofile (boschi a *Quercus ilex*), nelle quali, non di rado, accanto alla componente sempreverde è presente pure una componente caducifolia (*Fraxinus ornus*, ecc.). Il limite altimetrico inferiore attuale di queste formazioni (100-150 m) è legato all'azione umana di disboscamento per ottenere superfici per pascoli e coltivi (principalmente agrumi ed olivo).

- *Piano collinare*: si estende da 600 m fino a 1100 m di altitudine. La vegetazione zonale di questo piano è costituita da formazioni a latifoglie decidue termofile (*Quercus virgiliana*, *Q. cerris*, *Q. frainetto*, *Q. dalechampii*, *Ostrya carpinifolia*, *Acer obtusatum*, ecc.). E' in questo piano altitudinale che ha il baricentro la coltura del castagno.

- *Piano montano*: è situato oltre 1100 m di altitudine. E' caratterizzato da formazioni zonali a latifoglie decidue mesofile (*Fagus sylvatica*) ed a conifere (*Pinus laricio*). L'attività silvopastorale ha modificato notevolmente la vegetazione di questo piano, espandendo le aree a pascolo e favorendo in molte stazioni boschive il pino nei confronti del faggio.

I limiti altitudinali sopra riportati sono da considerarsi puramente indicativi in quanto la geomorfologia molto varia e diverse condizioni topoclimatiche possono determinare dislocazioni extrazonali di alcuni tipi di vegetazione.

Formazioni vegetali cartografate

La discussione delle formazioni vegetali segue l'ordine della legenda della carta annessa.

PIANO BASALE

Questo piano è caratterizzato da un notevole grado di antropizzazione, legato alla presenza dei centri urbani più estesi e popolati e di vaste superfici coltivate. La coltura più frequente è quella dell'olivo che dalla zona costiera si spinge sui rilievi interni fino a 500-600 m di quota. Nella piana di Rossano gli oliveti sono spesso associati agli agrumeti, mentre più a sud, sulle colline argillose del medio bacino del fiume Nicà, alla coltura degli agrumi si sostituisce quella della vite.

Formazioni a sclerofille sempreverdi termoxerofile

La lecceta

Le formazioni arboree del piano basale si presentano particolarmente frammentarie: si tratta per lo più di stadi di degradazione dei boschi a *Quercus ilex*, soggetti a diradamento tramite taglio ed incendio per aumentare le superfici a pascolo. La zona a NE di Rossano, comprendente i rilievi che salgono fino al santuario del Patire, è una tra le poche aree in cui la vegetazione a sclerofille sempreverdi si presenta con aspetti più maturi.

I rilievi fitosociologici eseguiti in questo tipo di vegetazione (Tab. 1) sono compresi in una fascia altitudinale che va da 140 m a 600 m di altitudine. Tra le specie che caratterizzano questa formazione boschiva prevalgono quelle stenomediterranee (*Quercus ilex*, *Phillyrea latifolia*, *Arbutus unedo*, *Asparagus acutifolius*, *Rubia peregrina*, *Selaginella denticulata*) seguite dalle eurimediterranee (*Fraxinus ornus*, *Celtis australis*, *Tamus communis*, *Cardamine graeca*), fra cui alcune con baricentro di distribuzione orientale. Notevole è pure la presenza nel sottobosco di numerose crittogame vascolari (*Asplenium trichomanes*, *Selaginella denticulata*, *Asplenium onopteris*, *Anogramma leptophylla*, *Polystichum* sp.). L'altezza dello strato arboreo non supera mai la decina di metri ed è sempre presente uno strato arbustivo ricco di specie e con elevati valori di copertura.

Dal punto di vista fitosociologico, le specie caratteristiche di *Quercetum ilicis* (*Arbutus unedo*, *Erica arborea*, *Phillyrea latifolia*, *Quercus ilex*, *Rubia peregrina*, *Smilax aspera*, ecc.) sono numerose e le fitocenosi in oggetto possono venir considerate una variante relativamente povera in elementi mediterraneo-orientali dell'associazione *Orno-Quercetum ilicis*. Barbero e Quezel (1976) hanno descritto un tipo di lecceta caratterizzato dalla presenza di *Arbutus unedo* e

Tab. 1 - Rilievi di lecceta. Le specie sporadiche sono riportate in appendice.

Relévs of evergreen sclerophyllous wood with Quercus ilex. The rare species are in the appendix.

Rilievo n°	1	2	3	4	5	6	7	8
Altitudine (m)	500	400	600	550	550	450	140	450
Esposizione	E	W	E	E	ENE	E	N	E
Inclinazione (°)	35	45	30	30	20	20	40	35
Superficie (m ²)	50	100	70	70	50	50	30	50
Roccia affiorante (%)	30	2	—	—	—	5	10	20
Copertura strato arboreo (%)	90	80	80	80	20	40	100	70
Strato arboreo:								
<i>Quercus ilex</i>	4	4	1	2	2	4	2	2
<i>Fraxinus ornus</i>	1	1	1	3		1		2
<i>Arbutus unedo</i>			1		+	1		
<i>Celtis australis</i>						+	2	2
Strato arbustivo:								
<i>Phillyrea latifolia</i>	2	+	1	+	1	1	2	2
<i>Quercus ilex</i>	3	+	+		2	+	2	+
<i>Fraxinus ornus</i>	r	+	+		+		+	1
<i>Erica arborea</i>	3	+	2	3	2		+	
<i>Cytisus villosus</i>		r	r				1	1
<i>Arbutus unedo</i>	1	r	r		3			
<i>Rubus</i> sp.		r	r			r		
<i>Calicotome villosa</i>				1	+			1
<i>Celtis australis</i>						r	2	2
Strato erbaceo:								
<i>Asplenium trichomanes</i>	+	+	r	+		+	+	+
<i>Selaginella denticulata</i>	1	r	+	+	+		1	1
<i>Asplenium onopteris</i>	1	+	r	+		+	+	+
<i>Rubia peregriana</i>		+	+	+		r	+	+
<i>Tamus communis</i>	r	+	r	r		r		
<i>Ruscus aculeatus</i>		+	+	+		1	+	
<i>Cyclamen</i> sp.	+	+	+	+		r		
<i>Asparagus acutifolius</i>	+	+	+	r				
<i>Teucrium siculum</i>			r	+			+	1
<i>Cardamine graeca</i>		r	+	+				
<i>Smilax aspera</i>		+	+	+				
<i>Anogramma leptophylla</i>		r		+		r		
<i>Geranium robertianum</i>	r		r		r			
<i>Carex distachya</i>		+	1		+			
<i>Hypericum barbatum</i>					r		+	1
<i>Brachypodium sylvaticum</i>					+		+	+
<i>Quercus ilex</i> pl.					1		+	+
<i>Polystichum</i> sp.				r			+	+

strettamente affine all' *Orno-Quercetum ilicis* su substrati acidi del Pindo e del Peloponneso, il quale è floristicamente molto simile alle fitocenosi qui rilevate. La povertà in elementi caratteristici mediterraneo-orientali di *Orno-Quercetum ilicis* in entrambi i tipi di fitocenosi è probabilmente di natura edafica: infatti le facies più ricche in specie caratteristiche di tale associazione sono quelle site su roccia calcarea, mentre in Sila Greca (come nelle zone rilevate dagli autori succitati) il substrato non è di natura carbonatica.

La macchia bassa

La macchia mediterranea di origine secondaria e dinamicamente collegata ai boschi sopra descritti è costituita da suffrutici, cespugli ed alberi di piccole dimensioni che non superano mai 2-3 m di altezza. Essa si presenta fitta su scarpate e versanti ad elevata inclinazione e diventa più rada su superfici pianeggianti soprattutto per effetto del pascolamento intensivo. Ove il degrado è maggiore, questi popolamenti sono quasi esclusivamente formati da *Calicotome villosa*, *Phillyrea latifolia*, *Asparagus acutifolius*, *Cistus monspeliensis*, *Cistus incanus*; nelle stazioni meno disturbate a queste specie si aggiungono *Rhamnus alaternus*, *Erica arborea*, *Pistacia lentiscus*, *Quercus ilex*.

La boscaglia dei greti delle fiumare

Lungo i corsi d'acqua e soprattutto sui greti delle fiumare nelle posizioni più umide cresce una vegetazione arbustiva caratterizzata da *Nerium oleander*, al quale si accompagnano poche altre specie legnose, fra cui *Vitex agnus-castus* e *Tamarix* sp. pl. Dal punto di vista fitosociologico, potrebbe trattarsi di resti di una boscaglia ripariale tipica delle zone semiaride, inquadrabile nella classe *Nerio-Tamaricetea* (Schneider e Sutter, 1982).

Formazioni erbacee

Il pascolo sulle argille

Sulle colline argillose della fascia basale, oltre ad estese superfici coltivate a cereali, si trovano vaste aree utilizzate a pascolo ricoperte da una comunità vegetale a *Hedysarum coronarium* e *Cynara cardunculus* (Bernardo *et al.*, 1990). La composizione floristica di questa fitocenosi risulta dalla prima colonna della tabella sintetica fitosociologica riportata in Tab. 2. In questa formazione erbacea, a differenza di altre sottoposte alle stesse condizioni climatiche ma su substrato non argilloso, è notevole la presenza di specie perenni (*Hedysarum coronarium*, *Cynara cardunculus*, *Scorzonera trachysperma*, *Carduus corymbosus*, *Plantago serraria*), fra cui molte geofite (*Bellevalia romana*, *Gladiolus italicus*, *Ophrys* sp.pl.) a crescita e fioritura precoce in corrispondenza della breve stagione primaverile. Numerose specie di questa comunità sono

Tab. 2 - Classi di frequenza delle specie nei pascoli dei tre piani altitudinali. 1=piano basale; 2=piano collinare; 3=piano montano.

Frequency classes of species in the pastures of the three altitudinal belts. 1=lowland belt; 2= hill belt; 3= mountain belt.

Gruppo di rilievi n°	1	2	3		
Numero di rilievi	9	8	11		
<i>Hedysarum coronarium</i>	V	I		<i>Hypericum perforatum</i>	IV
<i>Cichorium intybus</i>	V	II		<i>Gaudinia fragilis</i>	IV
<i>Sherardia arvensis</i>	V	II		<i>Trifolium arvense</i>	IV I
<i>Cynara cardunculus</i>	IV			<i>Hypochaeris radicata</i>	IV II
<i>Inula viscosa</i>	IV			<i>Cynosurus echinatus</i>	IV III
<i>Euphorbia exigua</i>	IV			<i>Chondrilla juncea</i>	III
<i>Linum strictum</i>	IV			<i>Silene gallica</i>	III
<i>Blackstonia perfoliata</i>	IV	I		<i>Tolpis virgata</i>	II
<i>Bellardia trixago</i>	IV	I		<i>Trifolium strictum</i>	II
<i>Anagallis arvensis</i>	IV	I		<i>Jasione echinata</i>	II
<i>Scorzonera trachysperma</i>	IV	I		<i>Centaurea jacea</i>	II
<i>Aegilops geniculata</i>	IV	II		<i>Echium italicum</i>	II
<i>Geranium dissectum</i>	IV	II	I	<i>Quercus ilex (plantula)</i>	II
<i>Mentha pulegium</i>	IV	III		<i>Origanum vulgare</i>	II
<i>Ononis alopecuroides</i>	III			<i>Legousia speculum-veneris</i>	II
<i>Sonchus oleraceus</i>	III			<i>Linaria arvensis</i>	II
<i>Plantago serraria</i>	III			<i>Trifolium nigrescens</i>	II
<i>Bellevalia romana</i>	III			<i>Oglistia gallica</i>	II
<i>Picris echioides</i>	III			<i>Silene conica</i>	II
<i>Pallenis spinosa</i>	III			<i>Poa annua</i>	II
<i>Hypochaeris achyrophorus</i>	III			<i>Micropus erectus</i>	II
<i>Foeniculum vulgare</i>	III			<i>Verbascum thapsus</i>	II
<i>Catananche lutea</i>	II			<i>Torilis arvensis</i>	II
<i>Carduus corymbosus</i>	II			<i>Convolvulus elegantissimus</i>	II
<i>Nigella damascena</i>	II			<i>Lathyrus sphaericus</i>	II
<i>Centaureum maritimum</i>	II			<i>Agrostis castellana</i>	III I
<i>Brachypodium distachyum</i>	II			<i>Vicia cracca</i>	III I
<i>Avena fatua</i>	II			<i>Carduus nutans</i>	II I
<i>Gladiolus italicus</i>	II			<i>Tolpis umbellata</i>	II I
<i>Polygala monspeliaca</i>	II			<i>Herniaria glabra</i>	II I
<i>Euphorbia helioscopia</i>	II			<i>Lotus angustissimus</i>	II I
<i>Crepis vesicaria</i>	II			<i>Sinapis pubescens</i>	II II
<i>Ophrys bertolonii</i>	II			<i>Cerastium semidecandrum</i>	II II
<i>Carex flacca</i>	III I			<i>Galium verum</i>	II III
<i>Reichardia picroides</i>	III I			<i>Anthyllis vulneraria subsp. maura</i>	I II
<i>Dactylis hispanica</i>	III II			<i>Trifolium ochroleucum</i>	I III
<i>Scabiosa maritima</i>	III II			<i>Plantago serpentina</i>	I III
<i>Parentucellia viscosa</i>	III II			<i>Anthemis montana</i>	II V
<i>Convolvulus arvensis</i>	III II			<i>Poa bulbosa</i>	II V
<i>Centaureum erythraea</i>	III III			<i>Phleum ambiguum</i>	I V
<i>Sonchus arvensis</i>	II I			<i>Festuca circummediterranea</i>	I V
<i>Tragopogon dubius</i>	II I			<i>Armeria canescens</i>	V
<i>Trifolium lappaceum</i>	II I			<i>Helianthemum nummularium</i>	V
<i>Vicia sativa</i>	II I			<i>Hieracium hypnorum</i>	V
<i>Catapodium rigidum</i>	II II			<i>Potentilla calabra</i>	V
<i>Trifolium pallidum</i>	II II			<i>Centaurea deusta</i>	III IV
<i>Chrysanthemum segetum</i>	II II			<i>Silene vulgaris</i>	II IV
<i>Hedypnois rhagadioloides</i>	II II			<i>Achillea nobilis s.l.</i>	II IV
<i>Scorpiurus muricatus</i>	II II			<i>Leopoldia comosa</i>	II I IV
<i>Medicago hispida</i>	II II			<i>Hypericum barbatum</i>	I IV
<i>Gastidium ventricosum</i>	II III			<i>Petrorhagia saxifraga</i>	I IV
<i>Oenanthe pimpinelloides</i>	I II			<i>Anthoxanthum odoratum</i>	I IV
<i>Anthemis arvensis</i>	I II			<i>Sedum tenuifolium</i>	IV
<i>Briza maxima</i>	I II			<i>Bunium bulbocastanum</i>	IV
<i>Echium plantagineum</i>	I II			<i>Thymus pulegioides</i>	IV
<i>Dipsacus fullonum</i>	I II			<i>Leontodon cichoraceus</i>	IV
<i>Potentilla reptans</i>	I II			<i>Rumex acetosella</i>	I III
<i>Prunella vulgaris</i>	I II			<i>Viola aethensis subsp. messanensis</i>	III
<i>Vicia bithynica</i>	I II			<i>Orchis sambucina</i>	III
<i>Avena barbata</i>	I II			<i>Cuscuta epithymum</i>	II
<i>Trifolium scabrum</i>	I II			<i>Silene alba</i>	II
<i>Medicago orbicularis</i>	I III			<i>Cirsium vallis-demonii</i>	II
<i>Lolium perenne</i>	III V	I		<i>Plantago lanceolata</i>	V II V
<i>Trifolium campestre</i>	II V			<i>Bellis perennis</i>	III III
<i>Aira cupaniana</i>		V	II	<i>Hypochaeris laetigata</i>	II III III
<i>Daucus carota</i>	III IV			<i>Dactylis glomerata</i>	II I III
<i>Bromus sterilis</i>	III IV			<i>Raphanus raphanistrum</i>	II I II
<i>Trifolium angustifolium</i>	I IV			<i>Sanguisorba minor</i>	II II I
<i>Vulpia ciliata</i>	IV			<i>Bromus hordeaceus</i>	II II I
				<i>Picris hieracioides</i>	I II I
				<i>Cynosurus cristatus</i>	I II I
				<i>Carlina vulgaris</i>	I II II

strettamente legate al substrato argilloso (*Hedysarum coronarium*, *Ononis alopecuroides*, *Carduus corymbosus*, *Catananche lutea*). Dove il carico di pascolamento è più pesante, le specie non appetite dal bestiame come *Cynara cardunculus* invadono gran parte della superficie del pascolo.

La "steppa" a sparto

Sempre su substrato argilloso, ma dove la pendenza maggiore dà luogo a fenomeni erosivi che possono portare alla formazione di veri e propri calanchi, si sviluppa una fitocenosi fisionomicamente simile alla vegetazione steppica dell'Africa settentrionale, in cui dominano *Lygeum spartum* e *Cymbopogon hirtus*. Si tratta di una fitocenosi floristicamente molto simile alle praterie di *Lygeum spartum* descritte per la Calabria da Gentile e Di Benedetto (1961) e da Schneider e Sutter (1982).

La gariga a elicriso

Negli alvei dei corsi d'acqua e sui depositi glareosi costieri in prossimità delle foci delle fiumare sono diffusi popolamenti misti ad erbe e suffrutici. Si tratta di una gariga caratterizzata dalla presenza di *Helichrysum italicum*, *Artemisia variabilis* e *Teucrium polium*, alle quali si accompagnano varie specie erbacee tipiche delle praterie xeriche mediterranee (*Echium plantagineum*, *Silene gallica*, *Hypochaeris radicata*).

La vegetazione del litorale sabbioso

L'ambiente costiero è in gran parte degradato e la vegetazione si presenta frammentaria e molto impoverita. La costa è ovunque bassa e, dove si ha accumulo di depositi sabbiosi, si sviluppa una vegetazione psammofila caratteristica molto discontinua rappresentata da popolamenti pionieri. Si tratta di fitocenosi ricche di specie caratteristiche della classe *Ammophiletea* (*Agropyron junceum*, *Eryngium maritimum*, *Echinophora spinosa*, *Cyperus kalli*), alle quali, spingendosi in direzione della zona afitoica, si accompagnano varie specie di *Cakiletea maritimae* (*Cakile maritima*, *Salsola kali*).

PIANO COLLINARE

Il paesaggio di questa fascia altitudinale è molto vario con contrasti accentuati dalla diversità dei substrati: ai rilievi prevalentemente argillosi tra i quali scorre il fiume Nicà si sostituiscono gradualmente procedendo da SE a NW substrati marnoso-argillosi, marnoso-calcarei e granitici. Tale variabilità del substrato e delle forme di erosione determinano una notevole diversificazione dei tipi di vegetazione.

La superficie boscata, pur essendo molto estesa, risulta essere molto degradata sia nella struttura che nella composizione, per l'uso contemporaneo che subisce sia per la produzione di legname sia come pascolo. Considerando che queste formazioni boschive sono ubicate nella zona collinare e nei medi bacini idrografici dove è più rilevante il dissesto idrogeologico, si intuisce quale importanza viene ad avere la ricostituzione e la salvaguardia di questo patrimonio forestale.

Formazioni a latifoglie decidue termofile

Il querceto deciduo

Per ciò che riguarda la vegetazione arborea, nel settore NW predominano boschi a *Quercus virgiliana* e *Quercus cerris*, mentre nella parte centrale e SE del territorio a *Quercus virgiliana* si sostituisce quasi completamente *Quercus frainetto*. Sui versanti a forte pendenza e con frequenti affioramenti rocciosi si trova nel sottobosco pure *Quercus ilex* in forma arbustiva, il quale, legato alle particolari condizioni edafiche e topoclimatiche, sale nelle stazioni rupestri fino oltre 1000 m di altitudine.

In base ai risultati dell'analisi nodale eseguita sui rilievi vegetazionali delle fitocenosi forestali della Sila Greca, questi boschi si differenziano dalle formazioni arboree degli altri due piani altitudinali per la presenza delle seguenti specie: *Cytisus villosus*, *Crepis leontodontoides*, *Teucrium siculum* e *Silene italica*. La prima di queste specie indica il carattere ancora tipicamente mediterraneo di questi boschi (in molti rilievi è presente ancora *Quercus ilex* nello strato arbustivo ed in quello erbaceo), *Crepis leontodontoides* e *Silene italica* sono specie caratteristiche dell'ordine *Quercetalia pubescentis* (boschi di caducifoglie termofile), mentre *Teucrium siculum* indica la presenza di un substrato a reazione acida. Va notato che le prime due specie citate hanno distribuzione mediterranea centro-occidentale, mentre *Teucrium siculum* è un endemismo dell'Italia meridionale appartenente al gruppo di *Teucrium scorodonia*, specie subatlantica. Quindi, come già osservato precedentemente nelle note fitogeografiche, già in questa fascia di vegetazione si manifesta l'infiltrazione di elementi occidentali in formazioni, la cui composizione floristica è ancora prevalentemente orientale.

Sulla carta allegata questi boschi sono stati inclusi in un unico tipo di formazione, in quanto presentano dal punto di vista fisionomico una struttura molto simile. L'analisi su base floristica ha permesso però di distinguere nell'ambito dei boschi in oggetto cinque diversi tipi vegetazionali. La loro distribuzione può essere individuata sulla carta in base alle indicazioni fornite dalle sigle delle specie arboree presenti. Va qui notato che l'indicazione *Quercus pubescens* riportata sulla carta va probabilmente corretta dal punto di vista

Tab. 3 - Classi di frequenza delle specie nei tipi di bosco deciduo del piano collinare. In colonna 1 è riportato il numero di rilievi, in cui le specie sono presenti.

Frequency classes of species in the deciduous wood-types of the hill belt. In column 1 the numbers are the occurrences of the species in the relevés.

Gruppo di rilievi n°	1	2	3	4	5	6
Numero di rilievi	3	9	6	9	15	5
Strato arboreo:						
<i>Acer mospessulanum</i>	2		I			
<i>Quercus ilex</i>	3	II			I	
<i>Quercus virgiliana</i>	3	V			I	
<i>Quercus cerris</i>	1	V	V	V	II	
<i>Acer obtusatum</i>	1	I	II	V		
<i>Pinus laricio</i>		I	II	V	I	I
<i>Ostrya carpinifolia</i>	1		I	III		
<i>Quercus dalechampii</i>				III		
<i>Quercus frainetto</i>		II			V	
<i>Castanea sativa</i>	2	II	II		I	V
Strato arbustivo:						
<i>Acer mospessulanum</i>	2		I			
<i>Fraxinus ornus</i>	2	I			I	
<i>Quercus virgiliana</i>	1	IV			I	
<i>Pirus pyraister</i>	1	I				I
<i>Cytisus scoparius</i>	1	I				I
<i>Spartium junceum</i>	1	I	I		I	
<i>Crataegus monogyna</i>	1	III	III		II	II
<i>Quercus ilex</i>	2	IV	II	III	II	
<i>Rubus</i> sp.	3	IV	V	V	II	IV
<i>Rosa</i> sp.	1	II	III	II	I	II
<i>Cytisus villosus</i>	3	V	V	V	IV	II
<i>Quercus cerris</i>		V	V	V	II	I
<i>Cistus salvifolius</i>		I			I	
<i>Prunus spinosa</i>		I	II		I	
<i>Prunus dulcis</i>		I	I		I	
<i>Sorbus torminalis</i>		I	I	II		
<i>Acer obtusatum</i>	1	I		V		I
<i>Pinus laricio</i>		I		V	I	
<i>Ilex aquifolium</i>		I		V		
<i>Ostrya carpinifolia</i>				V		
<i>Quercus dalechampii</i>				III		
<i>Quercus frainetto</i>		I			V	
<i>Erica arborea</i>	1				IV	
<i>Cistus incanus</i>		I			IV	I
<i>Calicotome villosa</i>		I			III	
<i>Castanea sativa</i>	2	I	I		I	V
Strato erbaceo:						
<i>Calamintha nepeta</i>	3					I
<i>Lamium maculatum</i>	2					II
<i>Quercus virgiliana</i>	3	III			I	
<i>Castanea sativa</i>	2	II		I	I	
<i>Ruscus aculeatus</i>	2	II		II	II	
<i>Campanula rapunculus</i>	1	II			I	
<i>Geranium robertianum</i>	1	II			I	I
<i>Briza maxima</i>	1	I			I	
<i>Viola alba</i>	2	I	II		III	
<i>Doronicum orientale</i>	2	II	I	II		I

<i>Hedera helix</i>	2	I	I	III		
<i>Quercus ilex</i>	1	III	I	II	II	
<i>Crepis leontodontoides</i>	3	V	V	I	III	III
<i>Silene italica</i>	3	IV	IV	III	I	II
<i>Luzula forsteri</i>	1	V	IV	IV	I	II
<i>Clinopodium vulgare</i>	2	IV	I	III	III	IV
<i>Fragaria vesca</i>	1	III	IV	II	I	IV
<i>Dactylis glomerata</i>	1	III	III	III	IV	IV
<i>Helianthemum sp.</i>		II				
<i>Vicia sp.</i>		II			I	
<i>Petrorhagia sp.</i>		II	I			
<i>Allium triquetrum</i>		II	I			
<i>Tolpis umbellata</i>		III	I	I		
<i>Reichardia picroides</i>		I	II	I		
<i>Cynosurus echinatus</i>		I	I	I		
<i>Fraxinus ornus</i>		I	I	I		
<i>Daphne laureola</i>		II	II	IV		
<i>Viola reichenbachiana</i>		II	I	II		
<i>Aremonia agrimonoides</i>		I	I	I	I	
<i>Achillea ligustica</i>		III	III	I	II	
<i>Teucrium siculum</i>		V	V	II	V	
<i>Festuca heterophylla</i>		II	III	V	I	
<i>Quercus cerris</i>		III	V	V	I	
<i>Anthoxanthum odoratum</i>		III	II	II	IV	
<i>Brachypodium sylvaticum</i>		III	IV	II	IV	I
<i>Pteridium aquilinum</i>		III	III	V	III	III
<i>Lathyrus venetus</i>		II	I	III	I	I
<i>Poa trivialis</i>		III	V		III	IV
<i>Knautia calycina</i>		I	II		II	
<i>Trifolium ochroleucum</i>		II	I		II	
<i>Veronica chamaedrys</i>		II	I	II		
<i>Scutellaria columnae</i>		II		II		
<i>Ranunculus repens</i>		II		II	I	
<i>Sedum tenuifolium</i>		II		II	I	
<i>Galium album</i>		II		I	I	
<i>Jasione montana</i>		II		I	I	
<i>Digitalis micrantha</i>		I		III	I	II
<i>Geranium versicolor</i>		II		II	I	III
<i>Melica uniflora</i>		II		II		II
<i>Tamus communis</i>			II		I	
<i>Helleborus foetidus</i>			II		II	I
<i>Carlina vulgaris</i>			II	I		
<i>Silene vulgaris</i>			II	II		I
<i>Acer obtusatum</i>	2		I	V		
<i>Pinus laricio</i>		I		III		
<i>Quercus dalechampii</i>				III		
<i>Ostrya carpinifolia</i>				II		
<i>Sesleria autumnalis</i>				II		
<i>Poa bulbosa</i>				II		
<i>Cyclamen sp.</i>				II		
<i>Epipactis helleborine</i>				II		
<i>Hypochoeris achyrophorus</i>				II		I
<i>Quercus frainetto</i>		I			II	
<i>Trifolium medium</i>					V	
<i>Oenanthe pimpinelloides</i>			I		IV	
<i>Galium lucidum</i>					II	I
<i>Potentilla micrantha</i>					II	
<i>Vicia incana</i>					II	
<i>Dorycnium hirsutum</i>					I	
<i>Hypericum barbatum</i>					I	II
<i>Galium odoratum</i>			I			II

sistematico, in quanto nelle zone dove sono stati eseguiti i rilievi fitosociologici è stata osservata unicamente *Quercus virgiliana*.

I cinque tipi di bosco individuati sono i seguenti:

Bosco a Quercus virgiliana-Acer monspessulanum (Tab.3, colonna 1): si tratta di boschi estremamente termofili, posti a ca. 600 m di altitudine, in diretto contatto con la lecceta della fascia sottostante. Gli strati arboreo ed arbustivo sono costituiti da varie specie di *Quercetalia pubescentis* (*Quercus virgiliana*, *Acer monspessulanum*, *Fraxinus ornus*), alle quali si accompagna *Quercus ilex*. Fra le specie boschive sono degne di nota *Pirus pyraeaster* (arbustivo), *Doronicum orientale*, *Clinopodium vulgare*, *Hedera helix* e *Ruscus aculeatus*. Sia nello strato arboreo che in quello erbaceo si nota una mescolanza di specie caratteristiche di *Quercetalia pubescentis* e di *Quercetea ilicis*.

Bosco a Quercus cerris (Tab.3, colonne 2, 3, 4): *Quercus cerris* è la specie arborea più diffusa in questa fascia altitudinale. Nella composizione floristica delle cerrete della Sila Greca mancano quasi del tutto le specie caratteristiche delle cerrete dell'Appennino Lucano centro-settentrionale (Aita *et al.*, 1977). Inoltre, a differenza di quanto avviene nel resto dell'Italia meridionale, il cerro e *Quercus frainetto* in Sila Greca non sono ben equilibrati nell'ambito delle singole fitocenosi: infatti si hanno formazioni ove predomina il cerro o formazioni ove predomina il farnetto. Sembra cioè che il ruolo preponderante giocato dal cerro (specie submediterranea nord-orientale) in tutto l'Appennino, probabilmente anche per motivi silvicolturali (Quezel e Bonin, 1980), si indebolisca in questa regione per dar spazio ad altre specie quercine a distribuzione principalmente sud-italico-balcanica (*Quercus frainetto* e *Quercus dalechampii*). Nonostante la povertà in specie delle cerrete in oggetto rispetto a quelle lucane, si possono notare anche nei boschi a *Quercus cerris* della Sila Greca le diverse tendenze messe in evidenza da Aita *et al.* (1977) per le cerrete lucane e cioè: una variante a carattere termofilo ricca in specie di *Quercetalia pubescentis* (v. *Bosco a Quercus cerris-Quercus virgiliana*); una variante a carattere mesofilo con prevalenza di elementi di *Fagetalia* (di cui è stato eseguito un solo rilievo); una variante nella quale un ruolo importante è giocato da *Ilex aquifolium* (v. *Bosco a Quercus cerris-Ostrya carpinifolia*). Va ricordato inoltre che è su questo tipo di boschi, con tutte le sue varianti, che l'uomo ha impiantato la coltura del castagno (Bonin e Gamisans, 1976).

Le specie che caratterizzano le cerrete della Sila Greca nei confronti degli altri tipi di bosco sono: *Festuca heterophylla*, *Daphne laureola* e *Luzula forsteri*. Si tratta di tre specie mesofile caratteristiche della classe *Querco-Fagetea*.

Bosco a Quercus cerris-Quercus virgiliana (Tab. 3, colonna 2): questa variante più termofila delle cerrete è caratterizzata dalla rarefazione delle specie mesofile succitate e dall'ingresso di alcune specie caratteristiche di *Quercetalia pubescentis* (*Quercus virgiliana*, *Achillea ligustica*, *Trifolium ochroleucum*, *Scutellaria*

columnae), alle quali si accompagnano varie specie a distribuzione stenomediterranea (*Allium triquetrum*, *Tolpis umbellata*, *Reichardia picroides*).

Bosco a *Quercus cerris*-*Ostrya carpinifolia* (Tab. 3, colonna 4): questo tipo di bosco è presente nella parte occidentale dell'area studiata a quote che si aggirano attorno a 1000 m. Si tratta di alcuni frammenti, dei quali i più estesi sono sul Cozzo del Pesco nella zona cartografata come querceto a *Quercus cerris* con grado di copertura di 40%-70%. Nello strato arboreo di questi boschi è sempre presente con esemplari di notevoli dimensioni *Ostrya carpinifolia*, alla quale si accompagnano altre specie arboree SE-Europee (*Quercus dalechampii*, *Acer obtusatum*) e *Pinus laricio*. In particolar modo *Quercus dalechampii* gioca un ruolo importante in questa formazione nei rilievi alle quote più elevate (al di sopra di 1000 m di altitudine). Questo tipo di bosco si sviluppa in stazioni a forte pendenza su un substrato a tessitura grossolana, ricco in scheletro e con basso contenuto di argilla, il quale deriva dal disfacimento di rocce cristalline. Va notato che su un substrato simile sono stati rilevati i boschi a *Quercus dalechampii* nel Rodope greco (Zoller et al., 1977). Nello strato arbustivo dominano specie che Pignatti (1979) indica come elementi della fascia colchica (*Ilex aquifolium*, *Daphne laureola*, *Hedera helix*). Nello strato erbaceo vanno ricordate *Digitalis micrantha*, endemismo appenninico, per l'elevato grado di presenza e, in alcuni rilievi, *Sesleria autumnalis* con elevati valori di copertura. E' interessante notare inoltre che in questo tipo di bosco sia *Crepis leontodontoides* che *Teucrium siculum* (caratteristici dei boschi misti di caducifoglie di questa fascia) si rarefanno.

Le stazioni qui rilevate sembrano corrispondere a quelle con elevata umidità dell'aria e bassa capacità idrica del suolo, considerate da Pignatti (1982b) quali stazioni primarie di *Ostrya carpinifolia* al margine meridionale del suo areale.

E' in questo tipo di bosco e nelle cerrete più mesofile che si trovano le specie considerate da Gentile (1969) quali caratteristiche dell'associazione *Aquifolio-Fagetum*, per cui è valida anche per la Sila Greca la seguente osservazione fatta da Aita et al. (1984) per la Lucania centro-settentrionale: "... le specie caratteristiche dell'*Aquifolio-Fagetum*, che dall'Appennino centrale verso nord appaiono più strettamente legate ai querceti caducifogli, anche nelle nostre zone, dove questi boschi sono ben più rappresentati, perdono gran parte del loro significato; anzi talune di esse sono addirittura assenti ... o più rare ..., proprio in faggeta". Quindi in Calabria l'associazione *Aquifolio-Fagetum* sembra essere unicamente presente dove si ha una forte riduzione della fascia dei querceti caducifogli per la discesa del faggio (p. es. sulla Catena Costiera tirrenica), dovuta probabilmente alla costante azione dei venti umidi occidentali. Al contrario, sul versante ionico la faggeta è situata a quote superiori, dove le condizioni termiche non permettono la presenza delle specie di *Aquifolio-Fagetum*.

Bosco a Quercus frainetto (Tab. 3, colonna 5): in questo tipo di bosco, a differenza delle altre formazioni a *Quercus frainetto* rilevate nell'Italia meridionale (Bonin e Gamisans, 1976; Aita et al., 1977), *Quercus cerris* è presente solo raramente. Il substrato è simile a quello del tipo di bosco precedente (derivato dal disfacimento di rocce acide), la quota invece è leggermente più bassa (compresa fra 700 m e 1000 m) ed il grado di inclinazione è generalmente minore. Il fatto che *Quercus frainetto* cresca a quote inferiori rispetto a *Quercus dalechampii* è stato notato anche per la Grecia (monte Tymphristos - Voliotis, 1976). Le specie che sono legate a questo tipo di fitocenosi sono: *Oenanthe pimpinelloides* (caratteristica di *Melitto-Quercion frainetto*), *Helleborus foetidus*, *Potentilla micrantha*, *Galium lucidum*, *Vicia incana* (tutte quattro caratteristiche o frequenti nei boschi dell'ordine *Quercetalia pubescentis*), *Erica arborea*, *Calicotome villosa*, *Cistus incanus* (si tratta di tre specie stenomediterranee), *Viola alba* e *Trifolium medium*. La ricchezza in specie mediterranee e soprattutto la presenza notevole di *Erica arborea* ricordano i raggruppamenti a *Quercus frainetto* descritti da Barbero e Quezel (1976) per il Peloponneso.

Il castagneto

Come già accennato in occasione della descrizione delle cerrete, il castagno (*Castanea sativa*) è abbastanza diffuso in questa fascia di vegetazione. Esso è presente, sia pur in modo discontinuo, da 600 m a 1000 m di altitudine e con esemplari sporadici, residui di antichi castagneti da frutto, scende fino a 300 m di quota nei dintorni di Rossano.

Il castagno forma boschi cedui monofitici anche di notevole estensione (valle del Trionto presso Longobucco), mentre di rado si rinviene all'interno dei querceti. Dal punto di vista floristico i castagneti costituiscono fitocenosi, nelle quali sono scarsamente rappresentate le specie caratteristiche delle cerrete (Tab. 3, colonna 6).

L'arbusteto da degrado boschivo

Vaste superfici della fascia collinare sono occupate da una formazione cespugliosa secondaria, la cui origine è determinata dal taglio dei boschi o dall'abbandono delle attività agro-pastorali. Queste situazioni, che rientrano nelle fasi dinamiche di ripresa naturale o di degrado del bosco, hanno una composizione floristica che risulta alquanto omogenea. Le specie dominanti sono *Spartium junceum*, *Cytisus scoparius*, *Crataegus monogyna*, *Prunus spinosa*, *Rosa* sp. e *Rubus* sp. Nelle situazioni di maggior degrado è presente una densa copertura di *Pteridium aquilinum*.

Formazioni erbacee

Si tratta di pascoli su substrati eterogenei a componente prevalentemente arenaceo-marnosa o calcareo-marnosa. Essi possono venir caratterizzati dalla presenza di *Cynosurus echinatus*, *C. cristatus*, *Briza maxima*, *Trifolium campestre*, *T. arvense* e *T. angustifolium*.

La composizione floristica di questi pascoli risulta da Tab. 2 (colonna 2). Molto ben rappresentate sono le terofite (*Vulpia ciliata*, *Trifolium arvense*, *Silene gallica*, *Anthemis arvensis*, *Echium plantagineum*, *Briza maxima*, *Trifolium strictum*). Molte specie sono strettamente legate al substrato sabbioso (*Silene conica*, *Echium plantagineum*, *Jasione echinata*) ed alcune sono caratteristiche delle formazioni effimere su suolo acido (*Tuberarietalia guttati*: p. es. *Silene gallica*, *Briza maxima*).

Dove il substrato diventa a tessitura più fine tendono a prevalere le specie perenni (*Lolium perenne*, *Agrostis castellana*, *Oenanthe pimpinelloides*), mentre dove il substrato è argilloso si formano praterie a *Hedysarum coronarium* e *Ononis alopecuroides* simili a quelle descritte per i dossi argillosi del piano basale.

PIANO MONTANO

Formazioni a latifoglie decidue mesofile

I rilievi eseguiti in tale tipo di bosco (Tab. 4) sono tutti al di sopra di 1300 m di altitudine. L'analisi nodale ha messo in evidenza, quali specie differenziali di queste fitocenosi, numerosi taxa caratteristici di *Fagetalia* e *Quercus-Fagetea* (*Fagus sylvatica*, *Galium odoratum*, *Orthilia secunda*, *Sanicula europaea*, *Poa sylvicola*, *Epipactis helleborine*, *Viola reichenbachiana*, *Aremonia agrimonoides*). Il carattere tipicamente europeo della florula di questi boschi si riscontra pure nelle specie non strettamente nemorali (di radura o di orlo boschivo), che, a causa della forte azione di disturbo, entrano a far parte del corteggio floristico di tali faggete (*Astragalus glycyphyllos*, *Galium album*, *Veronica officinalis*, *Silene vulgaris*). Notevole è la presenza pure di *Pinus laricio*, dovuta però in queste fitocenosi all'azione umana di rimboschimento. Spesso entrano in questi boschi, soprattutto nelle situazioni più aperte, alcune specie trasgressive delle cerrete (*Quercus cerris*, *Crepis leontodontoides*, *Festuca heterophylla*, *Achillea ligustica*) e dei pascoli (*Hieracium hoppeanum*, *Sedum tenuifolium*, *Trifolium pratense*, *Helianthemum nummularium*, ecc.).

I rilievi riportati in Tab. 4 sono relativi a stazioni molto disturbate: indice di ciò è la massiccia presenza di origine antropica di *Pinus laricio*. Probabilmente per tale motivo risultano rare le specie considerate da Gentile (1969) caratteristiche dell'alleanza meridionale-orientale *Geranio-Fagion*: infatti sono presenti unicamente *Geranium versicolor* e *Doronicum orientale*. Le faggete meno degradate sul versante ionico della Sila sono situate al di fuori dei limiti dell'area

Tab. 4 - Rilievi di faggeta. Le specie sporadiche sono riportate in appendice.
Relevés of beech wood. The rare species are in the appendix.

Rilievo n°	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25		
Altitudine (x10 m)	134	131	155	155	135	156	153	145	132	132	130	150	158	160	155	166	130	135	135	135	135	158	135	136	155		
Esposizione	SW	SW	SE	SW	SE	S	SW	N	SW	S	W	NW	NW	S	E	NE	NW	NW	SW	SW	NE	N	E	NE	E		
Inclinazione (°)	10	10	10	10	10	20	5	30	10	10	10	20	15	10	15	10	10	5	20	20	30	40	20	10	20		
Superficie (m²)	80	100	80	100	100	100	100	50	100	100	50	80	80	100	80	100	50	80	90	80	80	80	80	80	80		
Roccia affiorante (%)	10	10	10	5	-	-	5	-	-	-	5	-	-	5	5	-	-	-	-	-	-	10	-	-	-		
Cop.strato arboreo (%)	40	60	50	60	90	90	80	70	70	80	80	90	90	80	90	70	80	70	90	90	80	90	80	80	90		
Strato arboreo:																											
<i>Fagus sylvatica</i>				3	5	5						2	2	3	4	1			3	3	4	5	4	4	5		
<i>Pinus laricio</i>	3	3	2				4	3	3	4	4	3	3	2	2	4	4	3	3	3	2		3	3			
Strato arbustivo:																											
<i>Fagus sylvatica</i>	+	+	+	1	4	3	+	2	1	1	2	2	2	1	1	1	2	1	3	3	2	2	2	3	4		
<i>Rubus</i> sp.	+	+	+	+	+	+			+	+	+		+	+	+	+					+	+	+	+			
<i>Pinus laricio</i>			1					1	2	2	1	2	2	1		+	2	1	1	1	1			+			
<i>Cytisus scoparius</i>	3	1	1				+																				
<i>Prunus spinosa</i>							1	+		1							+										
<i>Crataegus monogyna</i>									1		+						+										
<i>Rosa</i> sp.				+	+				1	1	+	+	+														
<i>Cytisus villosus</i>								1				1	+					1	1	+							
Strato erbaceo:																											
<i>Brachypodium sylvaticum</i>	+		+	+	+		+		+	1	1	+	1	+	r	1	1	1	1	1	+		+	+	r		
<i>Peridium aquilinum</i>	2	+	+	+	+				2	3	1	+		+	+	+	1	1	+			+	1	+	+		
<i>Fagus sylvatica</i> pl.	1	+	+	+	1	+	+	+				1	+	+					+	+		+	+	+	1		
<i>Galium odoratum</i>				+	+	1			+			+	1	+	r	+		+		+	1	+	1	1	+		
<i>Pinus laricio</i> pl.	+	+	+			r	+	+	+	+	+	+	+	+		+		+					r		r		
<i>Ornithia secunda</i>	+			+	+	+						+					1		+	+			2	2	+		
<i>Sanicula europaea</i>	+		+	+					1	1	1		1		r		1		1	1							
<i>Poa sylvicola</i>					+	+	+				+			+				+			+	+	+	+	+		
<i>Vicia</i> sp.			r	r			+	+				r									+		r	+			
<i>Hieracium hoppeanum</i>	1	+	+																				+				
<i>Sedum tenuifolium</i>			+	+				+	+																		
<i>Trifolium pratense</i>	+						+																				
<i>Anthoxanthum odoratum</i>	+		+																								
<i>Helianthemum nummularium</i>		+	+							+																	
<i>Trifolium ochroleucum</i>	1	+																							r		
<i>Anthyllis vulneraria</i>	+	+	+																								
<i>Luzula forsteri</i>			+				+												+								
<i>Achillea ligustica</i>	+	+	+				1	+	1	+	r																
<i>Astragalus glycyphyllos</i>			+				+		1	+	1					+	1										
<i>Bellis perennis</i>	+	+	1				1		1	+	r						+	1									
<i>Crepis leontodontoides</i>							+	+	+													r					
<i>Orchis maculata</i>		+	+						+	+	+											r					
<i>Clinopodium vulgare</i>			+						+							r		+	+								
<i>Silene vulgaris</i>	+	+	+		+		+	+	+			r	r		+	+		+									
<i>Epipactis helleborine</i>	+	+		+	r	+			+					r	+						r						
<i>Veronica officinalis</i>				+		r				+	+				+	+	+		+								
<i>Hypochoeris achyrophorus</i>	+	+	1				1	+	1	+				r		+	+	+							r		
<i>Galium album</i>	+		1				+			r		+			+	+						+					
<i>Silene italica</i>			+	+		+	+	+				+	+		+	+		+				+					
<i>Quercus cerris</i> pl.			+					+				r	r														
<i>Festuca heterophylla</i>									1	1						+											
<i>Doronicum orientale</i>														+					+			+					
<i>Fragaria vesca</i>									1	+	1	+	+	+	+		1	+	+	1	+	+	+	+			
<i>Lathyrus venetus</i>												+	+	+	+	+	+	+	+	+	+	+	+	+			
<i>Dactylis glomerata</i>				+					+	+	+	+			r		+		+		+						
<i>Viola reichenbachiana</i>									+	+	+		+	+	+		1					+					
<i>Geranium versicolor</i>								+	1				+		+		1		+		+	+					
<i>Arenonia agrimonoides</i>				+										+		+	+				+	+	+				
<i>Mycelis muralis</i>																		+	+			+	+				
<i>Anthemis montana</i>	+						+											+	+			+	+		r		
<i>Galium lucidum</i>																+					+			r			
<i>Epilobium</i> sp.																	+					+	r				

della Comunità Montana e quindi non sono trattate nel presente lavoro. Per farsi un'idea su quale può essere la vegetazione potenziale di questa fascia, basti ricordare che esse sono caratterizzate dalla presenza di *Abies alba* e di *Campanula trachelium* e che corrispondono molto bene dal punto di vista floristico all'associazione *Asyneumat-Fagetum* di Gentile (1969).

Formazioni a conifere

Solo su un substrato costituito da sabbia grossolana derivata dal disfacimento dei graniti si trovano alcuni lembi di pineta probabilmente naturale, in cui entrano molti elementi tipici delle cerrete sottostanti. Queste formazioni su sabbia sono caratterizzate dalla presenza di *Hypochoeris laevigata* e *Cytisus sessilifolius*. Talvolta è presente pure *Quercus cerris*, mentre nelle schiarite sono frequenti *Astragalus calabrus* e *Chamaecytisus spinescens*. Questa vegetazione si presenta maggiormente sviluppata sull'altopiano silano, mentre nella zona studiata risulta molto frammentaria e ridotta. Va notato che i consorzi a *Pinus laricio* su sfaticcio di granito, con le numerose specie di pascolo e la quasi costante presenza di *Astragalus calabrus*, ricordano i rilievi eseguiti da Barbero e Quezel (1976) nei boschi a *Pinus pallasiana*-*Crataegus pycnoloba* del Peloponneso, dove *Astragalus parnassii* ha elevati valori di presenza. Già in Pignatti et al. (1980) sono state messe in evidenza le affinità orientali delle formazioni ad *Astragalus calabrus*, considerate uno stadio degradato della pineta a pino laricio, in netto contrasto con le affinità occidentali degli astragaleti di Sicilia, Sardegna e Corsica.

Formazioni erbacee

Sono costituite da pascoli e prati-pascoli xerici su substrato sabbioso. Dal punto di vista strutturale, la maggioranza delle specie sono perenni (ca. 90%, v. Codogno et al., 1987): soprattutto le camefite presentano elevati valori di frequenza (*Helianthemum nummularium*, *Thymus pulegioides*, *Sedum tenuifolium*, ecc.).

Dal punto di vista floristico, questa formazione può essere tipizzata come comunità a *Potentilla calabra*-*Armeria canescens* (Bernardo et al., 1990). Specie differenziali di questo tipo di fitocenosi sono: *Potentilla calabra*, *Armeria canescens*, *Festuca circummediterranea*, *Helianthemum nummularium*, *Hieracium hoppeanum*, *Sedum tenuifolium*, *Bunium bulbocastanum*, *Thymus pulegioides*, *Hypericum barbatum*, *Leontodon cichoraceus*, *Plantago serpentina*, *Petrorhagia saxifraga*, *Anthoxanthum odoratum* e *Phleum ambiguum*. Molte specie che concorrono a formare queste fitocenosi sono caratteristiche delle praterie su suolo acido sabbioso (*Sedo-Scleranthetea*: p. es. *Plantago serpentina*, *Petrorhagia saxifraga*, *Sedum tenuifolium*, ecc.). La stessa *Potentilla calabra* appartiene al gruppo di *Potentilla argentea*, la quale è specie caratteristica di *Sedo-Scleranthetea*.




Fig. 3 - Altitudine dei rilievi nei diversi tipi di bosco. Sono indicati il valore medio e la deviazione standard.

Altitude of the relevés in the different wood-types. The mean value and the standard deviation are reported.

- A: piano basale (lowland belt)
 B: piano collinare (hill belt)
 C: piano montano (mountain belt)

- 1: lecceta (evergreen sclerophyllous wood with *Quercus ilex*)
 2: bosco a *Quercus virgiliana*-*Acer monspessulanum*
 3: bosco a *Quercus cerris*
 4: bosco a *Quercus cerris*-*Quercus virgiliana*
 5: castagneto (chestnut wood)
 6: faggeta (beech wood)
 7: bosco a *Quercus frainetto*
 8: bosco a *Quercus cerris*-*Ostrya carpinifolia*

FORMAZIONI AZONALI

Una tipica formazione azonale è la vegetazione boschiva ripariale che accompagna i corsi d'acqua in tutte e tre le fasce di vegetazione individuate. Essa è costituita principalmente da *Alnus glutinosa*, *Populus nigra*, *Salix* sp. pl., *Sambucus nigra* ed è determinata dalla presenza di acqua superficiale.

Legate invece alle condizioni edafiche e morfologiche della fascia collinare e montana sono le formazioni ad *Alnus cordata*. Questa specie a rapido accrescimento si comporta da pioniera su substrati a forte erosione e a inclinazione elevata, dove si insedia in popolamenti radi o con esemplari sparsi.

Considerazioni conclusive

Dal confronto della zonazione verticale della vegetazione forestale della Sila Greca sintetizzata in Fig. 3 con i dati riportati in Horvat *et al.* (1974), Ozenda (1975) e Mayer (1984), risulta chiaro il parallelismo esistente fra la vegetazione dell'area studiata e quella della penisola balcanica meridionale (Pindo e Peloponneso) per le fasce di vegetazione a carattere prevalentemente mediterraneo. Con l'aumentare della quota queste somiglianze si indeboliscono sempre più e continuano a permanere solo in presenza di un substrato a tessitura grossolana derivato dal disfacimento di rocce granitiche. Sugli altri substrati prevalgono, con l'aumentare della quota, dapprima *Quercus cerris* e poi *Fagus sylvatica* in formazioni più o meno simili a quelle di tutto l'Appennino meridionale, però più povere di specie caratteristiche.

Appendice

Specie sporadiche nei rilievi fitosociologici:

Tab. 1 - Rilievi di Icceta.

In 2 rilievi - Strato arboreo: *Quercus virgiliana* (2,3); *Phillyrea latifolia* (3,4). Strato arbustivo: *Acer monspessulanum* (1,8); *Pistacia lentiscus* (2,6); *Quercus virgiliana* (3,8); *Cistus salvifolius* (5,6). Strato erbaceo: *Celtis australis* (1,8); *Sedum cepaea* (1,8); *Micromeria* sp. (1,8); *Acanthus* sp. (1,8); *Sesleria autumnalis* (2,4); *Pulicaria odora* (2,7); *Luzula forsteri* (4,5); *Quercus frainetto* (4,5); *Carex flacca* (4,5); *Allium triquetrum* (4,5); *Anemone apennina* (4,5); *Stellaria media* (4,5) *Crepis leontodontoides* (5,8); *Allium subhirsutum* (6,7).

In 1 rilievo - Strato arboreo: *Erica arborea* (4); *Quercus frainetto* (4); *Pirus pyraeaster* (7). Strato arbustivo: *Lonicera implexa* (3); *Prasium majus* (6); *Crataegus monogyna* (6); *Clematis flammula* (6); *Ligustrum vulgare* (7); *Cistus incanus* (7). Strato erbaceo: *Crucianella angustifolia* (1); *Arabis collina* (1); *Silene italica* (1); *Quercus virgiliana* (1); *Poa trivialis* (1); *Hypochoeris achyrophorus* (1); *Phillyrea latifolia* (1); *Dactylis glomerata* (1); *Fraxinus ornus* (1); *Limodorum*

abortivum (2); *Thalictrum calabricum* (4); *Melica uniflora* (5); *Alliaria petiolata* (5); *Symphytum tuberosum* (5); *Melittis melissophyllum* (5); *Trifolium pratense* (5); *Sonchus arvensis* (5); *Galium album* (5); *Anthoxanthum odoratum* (5); *Stipa bromoides* (6); *Clinopodium vulgare* (6); *Briza maxima* (8).

Tab. 4 - Rilievi di faggeta.

In 2 rilievi - Strato arbustivo: *Acer obtusatum* (16,23). Strato erbaceo: *Jasione montana* (1,3); *Hypericum barbatum* (1,7); *Centaurea* sp. (1,7); *Orchis sambucina* (3,4); *Malva moschata* (3,7); *Viola calcarata* (3,7); *Petrorhagia* sp. (3,7); *Cytisus sessilifolius* (6,8); *Rumex obtusifolius* (8,9); *Poa trivialis* (9,10); *Cruciata laevipes* (9,10); *Hieracium sylvaticum* (9,10); *Thalictrum calabricum* (10,17); *Digitalis micrantha* (11,14); *Geum urbanum* (18,19); *Reichardia picroides* (21,24).

In 1 rilievo - Strato arboreo: *Populus tremula* (10); *Alnus cordata* (21). Strato arbustivo: *Pirus pyraister* (5); *Acer pseudoplatanus* (11); *Castanea sativa* (11); *Alnus cordata* (21); *Quercus ilex* (24); *Abies alba* (25). Strato erbaceo: *Knautia calycina* (1); *Carlina vulgaris* (3); *Stellaria media* (4); *Cruciata glabra* (7); *Polygala vulgaris* (9); *Trifolium medium* (9); *Prunella vulgaris* (10); *Campanula trachelium* (13); *Cardamine kitaibelii* (14); *Castanea sativa* (16); *Abies alba* (17); *Scrophularia canina* (17); *Lysimachia* sp. (17); *Geranium robertianum* (18);

Ringraziamenti. Desideriamo ringraziare il Prof. D. Lausi e il Prof. P.L. Nimis (Trieste) per la revisione critica del manoscritto e la Dr.ssa G. Bolognini (Trieste) per l'assistenza prestata durante la stesura del testo.

Bibliografia

- Aita L., F. Corbetta e F. Orsino, 1977. Osservazioni fitosociologiche sulla vegetazione forestale dell'Appennino Lucano Centro-Settentrionale. 1. Le Cerrete. Arch. Bot. e Biogeogr. Ital., 53 (3-4): 97-130.
- Aita L., F. Corbetta e F. Orsino, 1984. Osservazioni fitosociologiche sulla vegetazione forestale dell'Appennino Lucano Centro-Settentrionale. 2. Le Faggete. Boll. Acc. Gioenia Sc. Nat., 17: 201-219.
- Barbero M. e P. Quezel, 1976. Les groupements forestiers de Grèce Centro-Méridionale. Ecol. Médit., 2: 3-86.
- Bernardo L., M. Codogno e D. Puntillo, 1990. The floristic composition of the Sila Greca pastures (Calabria, S-Italy). Studia Geobot., 10: 59-70.
- Bonin G., 1978. Contribution à la connaissance de la végétation des montagnes de l'Apennin centro-méridionale. Thèse doct., Marseille, St-Jérôme: 318 pp.
- Bonin G. e J. Gamisans, 1976. Contribution à l'étude des forêts de l'étage supraméditerranéen de l'Italie Méridionale. Doc. Phytosociol., 19-20: 74-88.
- Braun-Blanquet J., 1964. Pflanzensoziologie. Grundzüge der Vegetationskunde. 3a ed., Springer, Wien-New York.

- Češka A. e H. Roemer, 1971. *A computer program for identifying species-relevé groups in vegetation studies*. Vegetatio, 23: 255-276.
- Ciancio O., 1971. *Sul clima e sulla distribuzione altimetrica della vegetazione forestale in Calabria*. Ann. Ist. Sperim. Selv., 2: 321-372.
- Codogno M., L. Bernardo, S. Fascetti e D. Puntillo, 1987. *Correlation between ecological factors and structural features of pasture vegetation in the Sila Greca (Calabria, S-Italy)*. Giorn. Bot. Ital., 121 (3-4): 193-196.
- Codogno M. e D. Puntillo, 1987. *Contributions to the quantitative phytogeography of Calabria (S-Italy)*. 1. *The pasture vegetation of the Sila Greca*. Ann. Bot. (Roma), 45 (2): 55-63.
- Codogno M. e D. Puntillo, 1988. *Über die Veränderungen der Höhenstufenfolge in kleineren Bereichen des Mittelmeerraums (insbesondere innerhalb Kalabriens, S-Italien)*. Sauteria, 4: 141-148.
- Gentile S. e G. Di Benedetto, 1961. *Su alcune praterie a Lygeum spartum L. e su alcuni aspetti di vegetazione di terreni argillosi della Sicilia orientale e Calabria meridionale*. Delpinoa, n.s., 3: 67-151.
- Gentile S., 1969. *Sui faggeti dell'Italia meridionale*. Atti Ist. Bot. Univ. Pavia, ser. 6, 5: 207-306.
- Horvat I., V. Glavač e H. Ellenberg, 1974. *Vegetation Südosteuropas*. Geobotanica Selecta, 4, Fischer, Stuttgart.
- Mayer H., 1984. *Wälder Europas*. Fischer, Stuttgart-New York.
- Ozenda P., 1975. *Sur les étages de végétation dans les montagnes du bassin méditerranéen*. Doc. Cart. Ecolog., 16: 1-32.
- Pignatti S., 1979. *I piani di vegetazione in Italia*. Giorn. Bot. Ital., 113 (5-6): 411-428.
- Pignatti S., 1982a. *Flora d'Italia*. 3 voll., Edagricole, Bologna.
- Pignatti S., 1982b. *Die Beteiligung der Ostrya-Gesellschaften in den submediterranen Vegetationskomplexen*. Studia Geobot., 2: 211-216.
- Pignatti E. e S., P. Nimis e A. Avanzini (1980). *La vegetazione ad arbusti spinosi emisferici: Contributo alla interpretazione delle fasce di vegetazione delle alte montagne dell'Italia mediterranea*. C.N.R., Roma, Coll. P.F. "Promozione della Qualità dell'Ambiente", AQ/1/79.
- Puglisi S., G. Callegari, V. Cantore, F. Iovino, G. Pontecorvo e G. Trisorio Liuzzi, 1985. *Il ruolo delle sistemazioni idraulico forestali nel recupero di terre marginali in alcuni bacini torrentizi della Sila Greca*. In: AA.VV. *Marginalità e sviluppo*. C.N.R., I.P.R.A., Marra ed., Cosenza: 263-340.
- Quezel P. e G. Bonin, 1980. *Les forêts feuillues du pourtour méditerranéen. Constitution, Écologie, Situation actuelle, Perspectives*. Rev. For. Franc., 32 (3): 253-268.
- Schneider R., e R. Sutter, 1982. *Beitrag zur Flora und Vegetation Südkalabriens*. Phytocoenologia, 10 (3): 323-373.
- Thorntwaite C.W., 1948. *An approach toward a rational classification of climate*. Geogr. Rev., 38: 55-94.
- Voliotis D., 1976. *Die Gehölzvegetation und die Vegetationszonierung des Gebirgszuges Tymfristos-Oeta-Parnassos (Griechenland)*. Candollea, 31: 37-51.

- Wildi O. e L. Orlóci, 1983. *Management and Multivariate Analysis of Vegetation Data*. 2a ed., Berichte d. Eidg. Anst. f. forstl. Vers. (Birmensdorf), 215: 1-139.
- Zoller H., P.Geissler e N. Athanasiadis, 1977. *Beiträge zur Kenntnis der Wälder, Moos- und Flechtenassoziationen in den Gebirgen Nordgriechenlands*. Bauhinia, 6 (1): 215-255.

Indirizzi degli autori:

Dr.ssa Liliana Bernardo
Prof. Giuliano Cesca
Dr.ssa Simona Fascetti
Sig. Domenico Puntillo

Orto Botanico
Università della Calabria
I-87030 Roges di Rende (CS)

Dr. Michele Codogno

Dipartimento di Biologia
Università di Trieste
Via A. Valerio 32
I-34127 Trieste

Direttore responsabile: Prof. DUILIO LAUSI
Decreto del Tribunale di Trieste n. 735 del 30.5.1988
Fotocomposizione ed impaginazione:
STUDIO GAMMA - Trieste
Stampa:
TIP. ZENIT
Via Rivalto 1 - Trieste
Finito di stampare nel giugno 1991

CONTENTS

- Nimis P.L., Lazzarin A., Lazzarin G. & Gasparo D. Lichens and bioindicators of air pollution by SO₂ in the Veneto Region (NE Italy). 3
- Bernardo L., Cesca G., Codogno M., Fascetti S. & Puntillo D. Studio fitosociologico e cartografia della vegetazione della Sila Greca (Calabria). 77

CONTENTS

Nimis P.L., Lazzarin A., Lazzarin G. & Gasparo D. Lichens and bioindicators of air pollution by SO ₂ in the Veneto Region (NE Italy).	3
Bernardo L., Cesca G., Codogno M., Fascetti S. & Puntillo D. Studio fitosociologico e cartografia della vegetazione della Sila Greca (Calabria).	77